

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 260 521 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

27.11.2002 Bulletin 2002/48

(51) Int Cl.7: **C07K 16/42, C07K 16/46,
A61K 39/395, A61P 37/08**

(21) Application number: **02009009.8**

(22) Date of filing: **14.08.1992**

(84) Designated Contracting States:

**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE**

(30) Priority: **14.08.1991 US 744768**

07.05.1992 US 879495

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:

92918713.6 / 0 602 126

(71) Applicant: **Genentech Inc.**

South San Francisco, CA 94080-4990 (US)

(72) Inventors:

- **Jardieu, Paula M.**
Uttica, New York 13501 (US)

• **Presta, Leonard G.**

San Francisco, California 94109 (US)

(74) Representative: **Kiddle, Simon John et al**

**Mewburn Ellis,
York House,
23 Kingsway
London WC2B 6HP (GB)**

Remarks:

This application was filed on 23 - 04 - 2002 as a
divisional application to the application mentioned
under INID code 62.

(54) **Immunoglobulin variants for specific Fc epsilon receptors**

(57) Two classes of polypeptides derived from human IgE are described. One class binds selectively to the high affinity IgE receptor on mast cells and basophils, but not to the low affinity IgE receptor on B-cells, monocytes, eosinophils and platelets. The other class binds to the low affinity receptor, but not the high affinity receptor. The differential binding polypeptides of this in-

vention are useful in diagnostic procedures for IgE receptors or in the therapy of IgE-mediated disorders such as allergies. They also are useful in preparing antibodies capable of binding regions of IgE that participate in receptor binding.

BEST AVAILABLE COPY

EP 1 260 521 A2

Description

Background of the Invention

[0001] This invention relates to amino acid sequence variant anti-IgE antibodies and to polypeptides containing IgE sequences, especially IgE antagonists and to polypeptides capable of differential binding to Fc ϵ RI and Fc ϵ RII.

[0002] IgE is a member of the immunoglobulin family that mediates allergic responses such as asthma, food allergies, type 1 hypersensitivity and the familiar sinus inflammation suffered on a widespread basis. IgE is secreted by, and expressed on the surface of, B-cells. IgE synthesized by B-cells is anchored in the B-cell membrane by a transmembrane domain linked to the mature IgE sequence by a short membrane binding region. IgE also is bound to B-cells (and monocytes, eosinophils and platelets) through its Fc region to a low affinity IgE receptor (Fc ϵ RII, hereafter "FCEI"). Upon exposure of a mammal to an allergen, B-cells are clonally amplified which synthesize IgE that binds the allergen. This IgE in turn is released into the circulation by the B-cells where it is bound by B-cells (through the FCEI) and by mast cells and basophils through the so-called high affinity receptor (Fc ϵ RI, hereinafter "FCEH") found on the surface of the mast cells and basophils. Such mast cells and basophils are thereby sensitized for allergen. The next exposure to the allergen cross-links the Fc ϵ RI on these cells and thus activates their release of histamine and other factors which are responsible for clinical hypersensitivity and anaphylaxis.

[0003] The art has reported antibodies capable of binding to FCEI-bound IgE but not IgE located on FCEH (see for example WO 89/00138 and US patent 4,940,782). These antibodies are disclosed to be clinically advantageous because they bind to IgE found on B-cells or circulating free in the body, but do not bind to FCEH and thus will not activate mast cells or basophils. In addition, various amino acid sequence variants of immunoglobulins are known, e.g., "chimeric" and "humanized" antibodies (see, for example, U.S. Patent 4,816,567; WO 91/09968; EP 452,508; and WO 91/16927). Humanized antibodies are immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibody may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance as will be more further described *infra*. Also known *per se* are monovalent and bispecific antibodies.

[0004] It is generally understood that FCEH, like FCEI, binds to recognition site(s) in the IgE constant (Fc) domain. The IgE recognition site(s) for the two receptors are poorly defined, despite considerable effort in the past directed to the problem.

[0005] Over the past decade several studies have been undertaken to determine which portion of the IgE molecule is involved in binding to Fc ϵ RI and Fc ϵ RII. Essentially three approaches have been tried. First, peptides corresponding to specific portions of IgE sequence have been used as either competitive inhibitors of IgE-receptor binding (Burt *et al.*, *Eur. J. Immun.*, **17**:437-440 [1987]; Helm *et al.*, *Nature*, **331**:180-183 [1988]; Helm *et al.*, *Proc. Natl. Acad. Sci.*, **86**:9465-9469 [1989]; Vercelli *et al.*, *Nature*, **338**:649-651 [1989]; Nio *et al.*, *Peptide Chemistry*, 203-208 [1990]) or to elicit anti-IgE antibodies which would block IgE-receptor interaction (Burt *et al.*, *Molec. Immun.*, **24**:379-389 [1987]; Robertson *et al.*, *Molec. Immun.*, **25**:103-113 [1988]; Baniyash *et al.*, *Molec. Immun.*, **25**:705-711 [1988]). The most effective competitive peptide was a sequence that was 1000-fold less active than IgE (Burt *et al.*, *Eur. J. Immun.*, **17**:437-440 [1987]).

[0006] Helm *et al.*, *Proc. Natl. Acad. Sci.*, **86**:9465-9469 (1989) found that a peptide corresponding to IgE residues 329-409 blocked *in vivo* sensitization of human basophil granulocytes with human IgE antibodies. Further studies indicated that residues 395-409 were not essential for binding of the 329-409 peptide to Fc ϵ RI (Helm *et al.*, *Proc. Natl. Acad. Sci.*, **86**:9465-9469 [1989]). Note that the IgE sequence variants described below had the sequence of Padlan *et al.*, *Mol. Immun.*, **23**:1063 (1986), but that the immunoglobulin residue numbers used herein are those of Kabat *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, Md. 1987).

[0007] Vercelli *et al.*, *Nature*, **338**:649-651 (1989) used recombinant IgE peptides as well as anti-Fc ϵ monoclonal antibodies to investigate the B-cell (Fc ϵ RII) binding site of human IgE. They concluded that the Fc ϵ RII binding site is in Fc ϵ 3 near K399-V402.

[0008] Burt *et al.*, *Eur. J. Immun.*, **17**:437-440 (1987) investigated seven peptides for competition against rat IgE in binding to rat mast cells. Their most active peptide, p129, was 1000-fold less active than IgE. p129 corresponds to human sequence 439-453 which includes loop EF. Another of their peptides, p130, corresponding to residues 396-419 in the Fc ϵ 3 domain, had no activity.

[0009] Robertson *et al.*, *Molec. Immun.*, **25**:103-113 (1988) assessed IgE binding by sequence-directed antibodies induced by several synthetic peptides. They concluded that the sequence defined by their ϵ -peptide-4 (corresponding

to residues 446-460), was not significantly involved in receptor binding while the sequence defined by their ϵ -peptide-3 (corresponding to residues 387-401), was likely to be proximal to the IgE-receptor recognition site.

[0010] Nio *et al.*, *Peptide Chemistry*, 203-208 (1990) evaluated numerous peptides with respect to their ability to inhibit histamine release by human basophils *in vitro*. Only one peptide (peptide 2, Table 1), exhibited specific inhibition; this peptide encompassed residues 376-388. However, a larger peptide which incorporated this sequence (peptide 3, Table 1), had no inhibitory activity.

[0011] Second, mutations in IgE have been partially explored. Schwarzbaum *et al.*, *Eur. J. Immun.*, 19:1015-1023 [1989] (*supra*) found that a point mutant P404H (P442H by the numbering system used herein) had 2-fold reduced affinity for Fc ϵ RI on rat basophilic leukemia (RBL) cells, but the interpretation of this finding is controversial (Weetall *et al.*, *J. Immunol.*, 145:3849-3854 [1990]).

[0012] Third, chimeric molecules have been constructed. Human IgE does not bind to the murine receptor (Kulczycki Jr., *et al.*, *J. Exp. Med.*, 139:600-616 [1974]) while rodent IgE binds to the human receptor with a reduced affinity (Conrad, *et al.*, *J. Immun.*, 130:327-333 [1983]); human IgG1 does not bind to IgE receptors (Weetall *et al.*, *J. Immun.*, 145:3849-3854 [1990]). Based on these observations, several groups have constructed human-murine chimeras or human IgE-IgG chimeras. Weetall *et al.*, *J. Immun.*, 145:3849-3854 (1990) made a series of human IgG1-murine IgE chimeras and concluded that the Fc ϵ 2 and Fc ϵ 3 domains are involved in binding murine Fc ϵ RI while the Fc ϵ 4 domain is unlikely to be involved in binding to murine Fc ϵ RI (but may possibly be involved in binding to Fc ϵ RII). However, these conclusions are uncertain since they rest primarily on *lack* of binding by chimeras and three of five chimeras lacked some interchain disulfide bonds.

[0013] Nissim *et al.*, *EMBO J.*, 10:101-107 (1991) constructed a series of human-murine IgE chimeras and measured binding to RBL cells and concluded that the portion of IgE which binds with high affinity to the specialized Fc ϵ receptor on RBL cells could be assigned to Fc ϵ 3.

[0014] The results reported by these authors (e.g. Helm *et al.* and Burt *et al.*) are inconsistent. Further, in the case of anti-IgE antibodies it is difficult to eliminate the possibility of nonspecific blocking due to steric hindrance (Schwarzbaum *et al.*, *Eur. J. Immun.*, 19:1015-1023 [1989]). It is apparent that considerable confusion exists in the art as to the domains of IgE Fc which are involved in the binding of IgE to Fc ϵ H or in the maintenance of IgE conformation responsible for IgE binding to Fc ϵ H.

[0015] It is an object of this invention to identify polypeptides capable of differential binding to Fc ϵ EL and Fc ϵ H.

[0016] It is an object herein to determine an IgE domain which is implicated in Fc ϵ H receptor binding, but which is not involved in Fc ϵ EL receptor binding, and vice-versa.

[0017] It is another object herein to identify antagonists which are capable of inhibiting allergic responses, including antagonists that neutralize the Fc ϵ H or Fc ϵ EL receptor-binding domains of Fc ϵ , immunoglobulin analogues that bind Fc ϵ EL but do not bind Fc ϵ H, or that bind Fc ϵ H but not Fc ϵ EL and humanized anti-huIgE antibodies that bind to Fc ϵ EL-bound IgE but not to Fc ϵ H-bound IgE or which bind to IgE but do not induce histamine release or degranulation of mast cells.

[0018] It is another object to provide novel polypeptides for use in the assay of Fc ϵ receptors and for use as immunogens or for selecting anti-IgE antibodies.

Summary of the Invention

[0019] We have identified domains and specific residues of IgE which play an important role in binding IgE to its Fc ϵ EL and Fc ϵ H receptors, and based on this information we have designed polypeptides which remain capable of substantially binding to only one of these two receptors while being substantially incapable of binding to the other of the receptors. These polypeptides are referred to as differential binding polypeptides. In particular, differential binding polypeptides that bind Fc ϵ EL comprise IgE sequences in which one or more residues in loop EF or the β -strand D domain are varied, while Fc ϵ H-binding polypeptides comprise IgE sequences in which loop AB and/or β -strand B sequences are varied. Conversely, included herein are certain polypeptides comprising functional IgE loop EF and β -strand D domains but loop AB and/or β strand B domains having reduced functionality compared to wild-type, which bind differentially to Fc ϵ H, and polypeptides comprising functional loop AB and β -strand B domains but β -strand D and/or loop EF domains having reduced functionality compared to wild-type, which bind differentially to Fc ϵ EL.

[0020] The differential binding polypeptides of this invention are sufficiently homologous with the amino acid sequence of an IgE heavy chain that they retain the capability to bind Fc ϵ EL or Fc ϵ H, but are varied such that they exhibit reduced ability to bind to one of the two receptors as compared to native IgE. In various embodiments, the polypeptides of this invention additionally comprise cytotoxic polypeptides, detectable labels, conformation-restraining groups and/or amino acid sequences which are heterologous to IgE, e.g. sequences from receptors or immunoglobulins as further described below. In other embodiments, the differential binding polypeptides comprise IgE sequences in addition to the above-mentioned receptor binding domains, e.g., at least one variable domain capable of binding a predetermined antigen. In another embodiment, the differential binding polypeptide is an IgE variant which is monovalent for a pre-

determined antigen. In a still further embodiment, the differential binding polypeptide comprises an inactive IgE variable domain, i.e., one which is incapable of binding to any antigen, or which is devoid of a variable domain or functional CDR.

[0021] The differential binding polypeptides of this invention are useful in diagnostic procedures for IgE receptors or in the therapy of IgE-mediated disorders such as allergies. They also are useful in preparing antibodies capable of

binding regions of IgE that participate in receptor binding.

[0022] In an embodiment of this invention, variant anti-IgE antibodies are provided for use in diagnosis or for the therapy or prophylaxis of allergic and other IgE-mediated disorders. In particular embodiments of this invention anti-IgE variant antibodies are provided in which one or more human (recipient) light chain residues 4, 13, 19, 24, 29, 30, 33, 55, 57, 58, 78, 93, 94, or 104, or heavy chain residues 24, 37, 48, 49, 54, 57, 60, 61, 63, 65, 67, 69, 78, 82, 97 or 100 have been modified, preferably by substitution with the residue found in the corresponding position in the donor (generally murine) antibody. In preferred embodiments, the selected residues are light chain 13, 19, 58, 78, or 104, or heavy chain residues 48, 49, 60, 61, 63, 67, 69, 82 or 82c, and most preferably are heavy chain residues 60, 61 or light chain residue 78.

[0023] In other embodiments we provide antibodies which are capable of binding FcEL-bound IgE but which are substantially incapable of binding FcEH-bound IgE or inducing histamine release from mast cells or basophils, comprising a human Kabat CDR domain into which has been substituted a positionally analogous residue from a Kabat CDR domain of the murine anti-huIgE antibodies MAE11, MAE13, MAE15 or MAE17. Also provided herein are bispecific antibodies and IgE-monovalent antibodies; and humanized antibodies exhibiting an affinity for IgE which ranges from about 0.1 to 100 times that of MAE11.

Brief Description of the Figure

[0024] FIG. 1 depicts the sequence of human IgE FcE2 and FcE3 (SEQ. ID. 1). This particular sequence is from Padlan *et al.*, *Molec. Immun.*, 23:1063-1075 (1986). Residues are numbered according to Kabat (*supra*). "X" residues are included to align the Padlan IgE sequence with the Kabat numbering scheme. Sequences which were altered in preparing various IgE mutants are underlined; bold numbers below the lines denote the mutant number. β -strand residues are overlined; loop residues are defined by all residues intervening between two β -strands.

[0025] Fig. 2 depicts light and heavy chain sequences for MAE11 (SEQ.ID. 2 and 3), MAE13 (SEQ.ID. 4 and 5) and MAE15 (SEQ.ID. 6 and 7).

[0026] Fig. 3 depicts heavy and light chain sequences for HuMae11V1 (SEQ.ID 8 and 9).

[0027] Figs. 4a and 4b depicts the percent inhibition of IgE binding to FcEL and FcEH receptors, respectively, by murine monoclonal antibody Mae11 as well as 3 humanized variants (v1, v8 and v9).

[0028] Figs. 5a-5b compare the binding of the MAE11, MAE15 and MAE17 antibodies to various huIgE variants. MAE1 is provided as a control which binds to both B cells and mast cell-bound IgE. The mutants scheduled in the boxes in each figure are identified in Table 11.

Detailed Description of the Invention

[0029] The IgE analogue polypeptides of this invention contain an amino acid sequence which is homologous to that of a naturally occurring IgE and have the ability to bind specifically or differentially to FcEL or FcEH but, in varying degree, not to both. The degree of homology of such polypeptides to wild-type IgE is not critical since only enough IgE sequence needs to be retained to enable the IgE to bind differentially or specifically to one of the two receptors. In general, the polypeptides of this invention will be IgE Fc analogues and will be about from 80% to 99% homologous with a polypeptide sequence of a naturally occurring IgE heavy chain Fc region. Homology is determined by conventional methods in which all substitutions are considered to be nonhomologous (whether conservative or nonconservative) and in which the sequences are aligned to achieve maximal homology.

[0030] It will be understood that the IgE Fc residue numbers referred to herein are those of Kabat. In applying the residue teachings of this invention to other IgE Fc domains it will be necessary to compare the entire candidate sequence with the Fig. 1 sequence in order to align the residues and correlate the residue numbers. In addition, the identity of certain individual residues at any given Kabat site number may vary from IgE to IgE due to interspecies or allelic divergence. When for example it is stated that substitutions are introduced at residue R383 (human IgE) it will be understood that this means introducing a substitution at the same site in IgE even though this same site (in loop AB) may be located at a different residue number or may be represented in the parental or starting IgE by a residue which is different than that described by Kabat. However, for the sake of clarity and simplicity the residue numbers and identities of the Kabat human IgE heavy chain sequences will be used herein. Note that some Kabat residues were deleted in the Padlan sequence, in which case the Kabat numbering system is preserved by insertion of a spacer residue designated "X" (See Fig. 1).

[0031] Similarly, the Kabat system is used to designate immunoglobulin residues used in the preparation of variant,

e.g. humanized, anti-IgE immunoglobulins such as IgG, IgE, IgA or IgD. In preferred embodiments the recipient human immunoglobulin site is numbered in accord with Kabat subgroups III (V_H) consensus and κ subgroup I (V_L) consensus sequences. In order to determine which donor residues correspond to these Kabat consensus residues the sequences are maximally aligned, introducing gaps as necessary, using the variable domain cysteine residues as principal guideposts. Note that CDRs vary considerably from antibody to antibody (and by definition will not exhibit homology with the Kabat consensus sequences). Maximal alignment of framework residues (particularly the cysteines) frequently will require the insertion of "spacer" residues in the numbering system, to be used for the F_v region of the donor antibody. For example, the residue "29a" referred to *infra*. This represents an extra residue found in the murine donor antibody V_{H1} CDR for which a counterpart does not exist in the consensus sequence but whose insertion is needed to obtain maximal alignment of consensus and donor sequences. In practice, then, when a humanized antibody (ver. 1) is prepared from this donor it will contain V_{H1} with residue 29a.

[0032] The differential binding polypeptides of this invention typically contain about from 5 to 250 residues which are homologous to an IgE heavy chain Fc region, but ordinarily will contain about from 10 to 100 such residues. Usually, the IgE Fc3 and Fc4 regions will be present, with the Fc3 domain providing residues directly involved in receptor binding with Fc4 being present to ensure conformational integrity.

[0033] Generally, the IgE is human IgE, although animal IgE such as rat, murine, equine, bovine, feline or porcine IgE is included. As noted above, there will be variation in the residue identities and numbers for these IgEs compared to the Fig. 1 sequence.

[0034] FCEH and FCEL are respectively defined to be the high affinity IgE receptor (FCEH, Ishizaka *et al.*, *Immunochimistry*, 7:687-702 [1973]) found on mast cells or basophils, and the low affinity receptor (FCEHII, or CD23) found on cells involved in inflammation such as monocytes, eosinophils and platelets, as well as B-cells (Capron *et al.*, *Immun. Today*, 7:15-18 [1986]). FCEH and FCEL include alleles and predetermined amino acid sequence variants thereof which bind IgE. While FCEH contains several polypeptide chains, the binding of candidate polypeptides to its alpha chain is all that needs to be assayed since the alpha chain is the portion of FCEH which binds IgE.

[0035] Differential binding means that the polypeptide will bind to one of FCEL or FCEH to the extent of at least about 75% of the degree with which the homologous native IgE binds to that receptor, but will not bind to the other receptor at more than about 20% of the degree that the homologous IgE binds to the other receptor. Binding is determined by the assays of Example 3. Included within this invention are polypeptides that are capable of binding to one of the two receptors to a greater degree than native IgE.

FCEL-Specific Polypeptides

[0036] These polypeptides preferentially bind to the low affinity receptor. They typically contain FcE3 sequences in which residues within the β -strand D domain or loop EF have been substituted or deleted, and/or an additional residue inserted adjacent to one of such residues. For the purposes herein, the beta strand D domain extends from N418-X431 (Fig. 1, wherein X indicates a residue omitted from U266 IgE but found in the Kabat sequence) and loop EF extends from G444 to T453. A preferred FCEL-specific embodiment is mutant 6 (Table 6), in which the substitution of 4 residues within the human IgE heavy chain sequence K423-R428 substantially abolished FCEH binding. Other FCEL-specific embodiments comprising EF loop variants are mutants 85, 89 and the combination of 49, 51, 52, 83, 86 and 87. These sites (the D and EF domains) are believed to be the principal sites involved in binding IgE to FCEL. However, those skilled in the art will be able to routinely screen for optimal FCEL-specific polypeptides using the methods shown in the examples once it is understood that the beta-strand D and loop EF domains are the principal mutagenesis targets.

[0037] The preferred FCEL-specific polypeptide is one in which a residue has been substituted or deleted from within the β -strand D domain or loop EF, or both. For example, four residues were substituted in generating mutation 6, and any one or more of these substitutions may be responsible for the loss in FCEH binding while retaining FCEL binding. As for loop EF, which is involved in both FCEL and FCEH binding, it is desirable to screen both activities in order to select the FCEL-specific IgE variants. For example, mutant 85 (in which 9 IgE residues are substituted by analogously positioned IgG residues) is not detectably capable of binding to FCEH, but does bind to FCEL (see Table 11). On the other hand, conversion of site 444 from Gly to Leu abolishes binding to either receptor, while sites 447 and 452 are involved in binding to both receptors since changes at these locations prevent binding to FCEL but do not abolish FCEH binding.

Beta-Strand D Variants for FCEL Specificity

[0038] In general, D domain substitutions will be nonconservative, i.e., substituted residues generally will differ substantially from those found within the homologous native IgE in terms of charge, hydrophobicity or bulk. Typically, a maximum of 4 of 14 β -strand D domain residues are varied (and are usually residues 423, 424, 426 and/or 428), although typically any 1 to 5 of these residues are suitable for variation. In general, no more than 4 residues need to

be varied and optimally only one will be varied.

[0039] K423 and/or K426 are substituted with any of a residue selected from the group of Arg, His, Cys, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Ile, Leu, Ser, Thr, Asp, Glu, Gln and Asn, preferably Gly, Pro, Glu, Gln and Asp and most preferably Pro or Gln.

5 [0040] E424 and/or E425 are substituted with any of a residue selected from Asp, Asn, Gln, His, Lys, Arg, Cys, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu, Ile, Ser and Thr, preferably Arg, Lys, Pro, Gly and His and most preferably Arg.

[0041] R428 and/or R422 are substituted with Cys, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu, Ile, Ser, Thr, Asp, Glu, Asn, Gln, His, and Lys, preferably Cys, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu, Ile, Ser, Thr, Asp, Glu, Asn and Gln, and most preferably Tyr.

10 [0042] T421 is substituted with Cys, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu, Ile, Ser, Asp, Glu, Asn, Gln, His and Lys, preferably Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu, Ile, Asp, Glu, Asn, Gln, His and Lys, and most preferably Phe, Trp, Pro, Gly, Ala, Val, Leu and Ile.

[0043] S420 is substituted with Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu and Ile, and preferably Pro or Gly.

[0044] X429 is substituted with any other naturally occurring amino acid residue.

15 [0045] It is likely that optimal differential and FCEL binding activity will be achieved by a combination of mutations. Preferably, FCEH or FCEL binding, as the case may be, will be less than 10% of native homologous IgE, and optionally will range from undetectable to 3% of native homologous IgE, while binding to the other receptor ranges from at least about 75% of native homologous IgE to 90%, and preferably 95% to greater than 100%, e.g. 125%. The mutations should be as conservative as possible, i.e., involve as modest changes in hydrophobicity, charge or bulk as possible, yet still result in a polypeptide exhibiting these differential binding characteristics.

20 [0046] Any one or more of the β -strand D domain residues also may be deleted. Deletion of residues may possess the advantage of not introducing potentially immunogenic sites into the IgE analogue.

[0047] Examples of candidate β -strand D domain substitutional or deletional variants are set forth in the following Table 1. To determine the sequence of each variant, identify the residue for each variant number under each site. For example, the sequence of compound 19 comprises C388 E389 E390, etc.

TABLE 1

AA ¹	HulGE Site					
	423 K	424 E	425 E	426 K	427 Q	428 R
C	19	20		37		55
M	18	21		38		56
F	8, 80	22		39		57, 88
Y	7	23		40		4, 75, 83-84, 89, 97
W	6	24		41		58, 85
P	1, 74, 78-79, 89, 103	25, 97		42		59
G	5, 76-77	26		43		60
A	12, 98-99	27, 98, 100		44, 98, 101		61, 98, 102
V	13, 97	28		45		62
L	14, 81	29		46		63
I	15, 82	30		47		64
S	16	31		48		65, 103
T	17	32		49		66, 104, 105
D	9		79	50		67, 86

¹ Amino acid residue substituted into the analogue

TABLE 1 (continued)

AA ¹	HulGE Site					
	423 K	424 E	425 E	426 K	427 Q	428 R
E	9, 94	1, 3-19, 37-54, 55-72, 75, 89, 88, 90-93, 99, 101, 102, 105	1-72, 74, 76-78, 80-88, 93-94, 99, 100-105	51		68, 87
N	10	33		52, 79, 84	79	69
Q	11	34		3, 54, 75, 80, 82-83, 85-89, 103-104	1-72, 75, 77, 78, 80-95, 97-103, 105	70
H	83, 104	35, 78, 84		53		71
K	2-4, 20-72, 75, 85-88, 91-93, 100-102, 105	36, 77, 79, 94		1-2, 5-36, 55-72, 74, 76, 77-90, 91, 93-95, 97, 99, 100, 102, 105	104	72, 79
R	84	2, 74, 76, 80, 81 83, 85-87, 103-104	89			1-3, 5-54, 74, 76-78, 80-82, 90-92, 94, 99, 100-101
Δ^2	90, 95, 96	91, 95, 96	91, 96	92, 96	96	93, 95, 96

¹Amino acid residue substituted into the analogue²Signifies a deletion

[0048] Insertion of one or more extraneous residues adjacent to a residue within the β -strand D domain also falls within the scope of this invention. Typically, only one residue will be inserted, although from 2 to 4 or more residues can be inserted adjacent to any one site within the domain. Smaller numbers of inserted residues will be preferred in order to avoid the introduction of immunogenic sites. This, however, is merely a matter of choice. In general, insertions will be made at a single site, although insertions can be made adjacent to any two or more β -strand D domain residues.

[0049] Insertions typically are made between the following residues: 422 and 423, 423 and 424, 424 and 425, 425 and 426, 426 and 427, 427 and 428 and/or 428 and 429. The inserted residue or residues generally will exhibit charge, bulk or hydrophobicity character which is distinct from that of the flanking residues. For example, candidate insertions can be selected from the following Table 2.

TABLE 2

Insertion	β -strand D domain site ¹
Q	1, 2, 3, 4, 5, 7 or 8
D	1, 2, 3, 4, 5, 6 or 7
E	1, 2, 3, 4, 5, 6 or 7
F	1, 2, 3, 4, 5, 6 or 7
W	1, 2, 3, 4, 5, 6 or 7
P	1 or 2
K	2 or 3
R	2 or 3

¹422R - site 1 - 423K - site 2 - 424E - site 3 - 425E - site 4 - 426K - site 5 - 427Q - site 6 - 428R - site 7 - 429X y - site 8. Absence of a site indicates no insertion at that site.

1422R - site 1 - 423K - site 2 - 424E - site 3 - 3425E - site 4 - 426K - site 5 - 427Q - site 6 - 428R - site 7 - 429X y - site 8. Absence of a site indicates no insertion at that site.

[0051] The FCEL-specific polypeptides are provided as linear or conformationally restrained polypeptides. Conformational restraint is accomplished by cross-linking the polypeptide, preferably at the N- and C- termini so as to produce a cyclic structure. In preferred embodiments the cyclic forms have the following structure:

O=C1N(R3)C(=O)C(R2)(R4)N1C(=O)C(R6)(R5)X

Chemical structure diagram showing a substituted amide. The structure includes a carbonyl group (C=O) connected to a nitrogen atom (N). The nitrogen atom is substituted with R₂ and R₃. The carbonyl carbon is substituted with R₄ and R₅. The nitrogen atom is also substituted with R₆ and R₇. The carbonyl carbon is substituted with R₈ and R₉. The structure is labeled with (a3-a11)(a12)(a13)(a14)(a15)Pro(a17)(a18)(a19) and a COR₁ group.

(a3-a11) is a bond or the sequence -R373 -F381; a12 and a18 are hydrophobic amino

	acid residues; a13 and a14 are basic amino acid residues; and a15, a17 and a19 are hydrophilic amino acid residues;
R ₁	is selected from
5	(a) hydroxy,
	(b) C ₁ -C ₈ alkoxy,
	(c) C ₃ -C ₁₂ alkenoxy,
	(d) C ₆ -C ₁₂ aryloxy,
	(e) acylamino-C ₁ -C ₈ -alkoxy
10	(f) pivaloyloxyethoxy,
	(g) C ₆ -C ₁₂ aryl-C ₁ -C ₈ -alkoxy where the aryl group is unsubstituted or substituted with one or more of the groups nitro, halo, C ₁ -C ₄ -alkoxy, and amino;
	(h) hydroxy substituted C ₂ -C ₈ substituted alkoxy; and
	(i) dihydroxy substituted C ₃ -C ₈ alkoxy;
15	
R ₂ , R ₃ , R ₅ , R ₇ , R ₈	are the same or different and are selected from
	(a) hydrogen,
20	(b) C ₆ -C ₁₂ aryl where the aryl group is unsubstituted or substituted by one or more of the groups nitro, hydroxy, halo, C ₁ -C ₈ alkyl, halo-C ₁ -C ₈ alkyl, C ₁ -C ₈ alkoxy, amino, phenyl, acetamido, benzamido, di-C ₁ -C ₈ alkylamino, C ₆ -C ₁₂ aroyl, C ₁ -C ₈ alkanoyl, and hydroxy substituted C ₁ -C ₈ alkyl,
	(c) C ₁ -C ₁₂ alkyl or alkenyl; C ₃ -C ₁₀ cycloalkyl or C ₃ -C ₁₂ substituted with any of halo, C ₁ -C ₈ alkoxy, C ₆ -C ₁₂ aryloxy, hydroxy, amino, acetamido, C ₁ -C ₈ alkylamino, carboxy or carboxamide;
25	
R ₂ and R ₃ , R ₅ and R ₆ , or R ₇ and R ₈	may optionally and independently be joined together to form a carbocyclic or heterocyclic ring of from four to seven atoms where the heteroatoms are selected from O, S, or NR ₁₀ where R ₁₀ is selected from
30	hydrogen, C ₁ -C ₈ -alkyl, C ₂ -C ₈ -alkenyl, C ₆ -C ₁₂ -aryl, C ₃ -C ₁₀ cycloalkyl, C ₆ -C ₁₂ -aryl-C ₁ -C ₈ -alkyl, C ₁ -C ₈ -alkanoyl, and C ₆ -C ₁₂ aroyl,
R ₄	is selected from
	hydrogen, C ₁ -C ₈ -alkyl, C ₂ -C ₈ -alkenyl, C ₆ -C ₁₂ -aryl, C ₃ -C ₁₀ cycloalkyl, C ₆ -C ₁₂ -aryl-C ₁ -C ₈ -alkyl, C ₁ -C ₈ -alkanoyl, and C ₆ -C ₁₂ aroyl;
35	R ₂ or R ₃ may be optionally joined with R ₄ to form a piperidine, pyrrolidine or thiazolidine ring;
X	is selected from
	an O or S atom,
40	NR ₉ wherein R ₉ is hydrogen, C ₁ -C ₈ -alkyl, C ₃ -C ₈ -alkenyl, C ₃ -C ₁₀ cycloalkyl, C ₆ -C ₁₂ -aryl, C ₆ -C ₁₂ -aryl-C ₁ -C ₈ -alkyl, C ₁ -C ₈ -alkanoyl, or C ₆ -C ₁₂ aroyl;
	C ₆ -C ₁₂ aryl,
	C ₁ -C ₈ alkanoyl, and
45	(CH ₂) _k where k is an integer from 0 to 5; and
	pharmaceutically acceptable salts thereof.
[0052]	As used herein and unless specified otherwise: alkyl and alkenyl denote straight or branched, saturated or unsaturated hydrocarbon chains, respectively; C ₆ -C ₁₂ aryl groups denote unsubstituted aromatic rings or fused aromatic rings such as, for example, phenyl or naphthyl; halo denotes F, Cl, Br, or I atoms; alkoxy denotes an alkyl group bonded through O to the indicated site. Examples of C ₁ -C ₈ alkyl or C ₂ -C ₈ alkenyl groups include methyl, ethyl, propyl, isopropyl, butyl, t-butyl, pentyl, isopentyl, hexyl, vinyl, allyl, butenyl and the like; examples of C ₃ -C ₁₀ -cycloalkyl groups include cyclopropyl, cyclopentyl, cyclohexyl, and the like; heterocyclic rings include but are not limited to pyridyl, thienyl, furyl, indolyl, benzthienyl, imidazolyl, thiazolyl, quinoliny and isoquinoliny. Hydrophobic amino acid residues include naturally occurring or synthetic residues having hydrophobic side chains, e.g. Phe, Leu, Ile, Val, Norleu, and the like.
50	Hydrophilic amino acid residues include naturally occurring or synthetic residues having charged or uncharged hydrophilic side chains, e.g. ornithine, Ser, Thr, Tyr, His, Asp, Glu, Lys and Arg. Preferably a15, a17 and a19 are unchanged and bear normal, secondary or tertiary mono or di-hydroxy substituted alkyl side chains. Basic residues have guanidino or amino-substituted side chains for the most part.
55	

[0053] The AB-B domain and/or loop EF - containing, FCEL-specific polypeptides of this invention optionally are associated with other substances or are fused to additional polypeptide sequences. The polypeptides generally contain only IgE-homologous sequences, although they also or alternatively are labelled for diagnostic use (employing enzymes, radioisotopes, biotin or avidin, stable free radicals, and chemiluminescent or fluorescent moieties in conventional fashion). Also the polypeptides are fused to non-IgE polypeptides such as cytotoxic or immunosuppressive polypeptides, to other IgE polypeptides (e.g. Fv regions), or to polypeptides capable of binding to a predetermined ligand or antigen.

[0054] Cytotoxic polypeptides include IgG Fc effector sequences and polypeptide toxins such as diphtheria toxin or ricin A chain (U.S. Patents 4,714,749 and 4,861,579). A preferred fusion is one in which the FCEL-specific sequence (such as that of the Fce3 - Fce4 sequence of mutant 6) is fused at its N-terminus (i.e., at approximately D360) to the C-terminus of an immunoglobulin, or an immunoglobulin fragment terminating at the C-terminus of IgG Fc γ 2 or IgG Fc γ 3. Alternatively the FCEL specific polypeptide is fused to an effector IgG sequence in place of one or both of the IgG Fv domains in analogous fashion to known immunoadhesins.

[0055] The polypeptides herein optionally are fused to polypeptides which are capable of binding a predetermined antigen or ligand. Generally, these additional polypeptides will be IgE or other immunoglobulin Fv domains, although they optionally are heterologous polypeptides such as receptor extracellular domains (produced in the known fashion of immunoadhesions, e.g. as has been accomplished with CD4). Immunoglobulin sequences fused to the FCEL-specific polypeptides herein include Fc or variable sequences of the heavy chains of IgG1, IgG2, IgG3, IgG4, IgE, IgM, IgD or IgA. Any FCEL-specific heavy chain fusion optionally is disulfide bonded in the ordinary fashion to heavy chains having the same specificity (thereby forming homopolymers) or to different heavy chains (thereby forming heteropolymers), including different heavy chains having specificity for a different antigen. Such heteropolymeric heavy chains include heavy chains which are not FCEL-specific, e.g., these comprise native IgE sequences which bind to FCEL and FCEH in the ordinary fashion, or the heavy chains optionally include at least one heavy chain that is FCEL specific and at least one that is FCEH specific. Heteropolymeric heavy chains also may include the heavy chains of non-IgE immunoglobulins, e.g., IgG, IgD, IgM and the like. In addition, the heavy chain hetero- or homopolymers optionally are disulfide bonded to light chains in the fashion of native immunoglobulins so as to cooperatively bind to predetermined antigen in the usual way. Unless the heteropolymeric heavy chains comprise IgM heavy chains they generally will be heterodimeric.

[0056] In some embodiments, immunoglobulins comprising a FCEL-specific polypeptide will also comprise an immunoglobulin variable region, preferably (if at all) an IgE Fv domain. The antigenic specificity of the variable region may vary widely, including those which bind haptens, or which bind polypeptides or proteins from human, animal, plant, fungal, bacterial or insect sources. The specificity may be unknown or the variable region may have the ability to bind to a predetermined antigen. If the immunoglobulin is to have a functional variable domain (as opposed to a deleted Fv in the case of Fce3 or Fce4 fragments) it is preferred that it have a known antigenic specificity. Antigenic specificity may include the ability to bind antigens associated with a cytotoxic or immune response, particularly lymphoid cell antigens such as CD3 or CD8, integrins, B-cell surface antigens, helper or suppressor cell surface antigens, or epitopes located in the variable region of effector subtypes of IgG. FCEL-specific Fc domains also are usefully employed in combination with Fv domains capable of binding a particular allergen to which a patient is allergic. These generally are human IgEs directed against allergens and which contain an FCEL-specific Fc domain. Alternatively, the immunoglobulin specificity is directed against the Fc region of effector subtypes of IgG, in this case however it being preferable that the FCEL-specific polypeptide not suppress complement binding or ADCC functions of the IgG.

[0057] The polypeptides of this invention that contain antigen or ligand binding capability contain one or more sites capable of binding to the antigen or ligand. For example, the polypeptides herein comprise one or more IgE or other immunoglobulin Fv domain to produce monovalent or polyvalent immunoglobulins. For the most part such polypeptides will be monovalent for antigen or ligand, as in the case when the immunoglobulin comprises a heavy-light chain pair that has a deleted or inactivated Fv or CDR so as to not be able to bind to antigen. Alternatively, they will be bivalent in the predominant instance, and will be monospecific or bispecific.

[0058] In another embodiment, FCEL-specific polypeptides are covalently bound to a cytotoxic agent. For example, the polypeptide ricin D toxin isolated from the *Ricinus communis* plant is bound to the carboxy terminus of the Fc domain, either by chemical means or, most preferably, by production of a fusion protein using standard recombinant DNA methods. This provides a means to selectively deliver the toxin only to cells expressing FCEL on their surfaces.

[0059] The FCEL-specific polypeptides need only contain so much of the IgE Fc ϵ sequence as is required to substantially maintain FCEL binding. This is readily determinable by synthesizing or expressing the product and determining its activity. In general, the entire IgE sequence extending from Fc ϵ 2 - Fc ϵ 4 can be used, although fragments containing only Fc ϵ 3 and Fc ϵ 4 are generally satisfactory.

[0060] In general the immunoglobulin sequences and the FCEL-specific sequence will be derived from the same species which is to be treated with the IgE analogue. Preferably, the immunoglobulin sequences are human.

[0061] The FCEL-specific polypeptides of this invention (when employed as such without fusion to non-IgE sequenc-

es) exclude the linear polypeptide sequences disclosed by Nio *et al.* (*supra*), as well as other prior art polypeptides which include the native IgE AB-B domain or loop EF (Burt *et al.*, *supra*).

FCEH-Specific Polypeptides

[0062] These polypeptides are amino acid sequence variants of IgE or its fragments in which a residue within the AB-B or loop EF domains have been deleted, substituted or another residue inserted so that the AB-B or loop EF domains are no longer capable of binding to FCEL, and which contain sufficient beta strand D sequence and (optionally) loop EF sequence to bind to the high affinity receptor. As disclosed above, the AB-B and loop EF domains have been implicated in binding to FCEL since mutations in these domains have a serious impact on the binding of the IgE variants to the low affinity receptor. In particular, mutations in loop EF or the C-terminal half of the AB loop and in the N-terminal half of beta strand B produce a divergence in IgE FCEL/FCEH specificity wherein the variant continues to bind to the high affinity receptor but largely fails to bind to the low affinity receptor. In addition, we have found that the IgE loop EF and the heavy chain beta strand D domains participate in binding to the high affinity receptor. Therefore, FCEH-specific differential binding polypeptides will comprise at least the FCEH-binding sequence of beta strand D and preferably also will contain a variant AB-B or loop EF domain sequence that binds substantially only to FCEH.

[0063] In preferred embodiments amino acid sequence variation is introduced into the low affinity receptor binding functionality of the AB-B or loop EF domains. Preferably, one or more of residues I382, R383, K384, S385, T387, I388, T389, C390, R446, D447, W448, I449, E150, G151, E152 or T153 are varied, although modifications optionally are introduced into loop AB N-terminal to the designated loop AB residues. Only one of R383, K384, S385, T387, T-389, or R446 - T453 need be mutated, although it is preferable to vary 1, 2 or 3 residues from each domain.

[0064] When substituted at all, I382 and/or I388 generally are independently substituted with Asn, Gln, Leu, Val, His, Lys, Arg, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Ser, Thr, Asp or Glu, preferably Trp, Pro, Gly, Ser, Thr, Asp or Glu. Ordinarily these two residues are not modified.

[0065] R383 typically is substituted with Cys, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu, Ile, Ser, Thr, Asp, Glu, Asn, Gln, His, or Lys, preferably Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu, Ile, Ser, Thr, Asp, Glu, Asn or Gln and most preferably Ala, Glu, Asp or Ser.

[0066] K384 typically is substituted with Arg, His, Cys, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Ile, Leu, Ser, Thr, Asp, Glu, Gln and Asn, preferably Ala, Gly, Pro, Glu, Gln or Asp and most preferably Ala, Glu or Asp.

[0067] S385 is substituted with Asp, Asn, Gln, His, Lys, Arg, Cys, Met, Phe, Tyr, Trp, Pro, Gly, Ala, Val, Leu, Ile, Glu and Thr, preferably Ala, Tyr, Val, Ile, Leu, Phe, Arg, Lys and His and most preferably Ala, Val, Ile, Leu, Phe and Tyr.

[0068] When substituted, P386 usually is substituted by Gly, Ala, Cys, Val, Leu, Ile, Ser, Thr, Asp, Glu, Asn, Gln, His, Lys, Arg, Phe, Tyr, or Trp, and preferably Gly, Ala, Ser, Thr, Asp, Glu, Asn, Gln, His, Lys, Arg or Trp. Ordinarily, P386 is not modified.

[0069] T387 and/or T389 generally are independently substituted by Gly, Ala, Val, Leu, Ile, Ser, Asp, Pro, Glu, Asn, Gln, His, Lys, Arg, Cys, Phe, Tyr and Trp, preferably Gly, Ala, Val, Leu, Ile, Asp, Glu, Asn, Gln, His, Lys, Arg, Phe, Tyr and Trp, and most preferably Ala.

[0070] C390 ordinarily is not substituted except when employed as a component of a cyclizing group as shown in Formula I.

[0071] The differential FCEH-binding polypeptides of this invention will comprise the sequence of functional FCEH-binding beta strand D and loop EF domains, as-defined above. In general, it is expected that the functional domains need not contain all of the beta strand D or loop EF domain residues. However, any modifications of the beta strand D domain residues will need to be conservative, if made at all, in order to preserve FCEH binding. Since loop EF is involved in both FCEL and FCEH binding, it likely will be necessary to screen these variants in order to determine their activity as shown in Example 5. However, a number of loop EF mutants already have been identified that substantially abolish FCEL binding without apparently interfering with FCEH binding, e.g. mutants 50 and 52. Thus, loop EF variants may belong in either the FCEL or FCEH specific category, or may equally affect binding to each receptor.

[0072] A particularly preferred embodiment of a FCEH-specific polypeptide is one which contains a beta strand D domain together with additional C-terminal sequence. The sequence of this embodiment extends from about T421 to about T440. Generally, the N-terminus of this embodiment is S420 or T421, while the C-terminus is T440, L441 or P442. In addition, one or more residues extraneous to this sequence are fused to its N- or C-termini. These extraneous residues are particularly useful in forming covalent or noncovalent bonds between the N- and C-termini of this polypeptide. The N- and/or C-termini preferably are covalently bonded through a side chain of a residue or through the polypeptide backbone. For example, cysteine residues are fused to the N- and C-termini and, upon oxidation, a polypeptide having a terminal disulfide bond is formed which joins the terminal ends of the polypeptide, thereby conformationally restraining the polypeptide. Alternatively, the alpha amino group of the polypeptide (or that of an extraneous N-terminally located residue) is covalently bonded to the sulfur atom of an extraneous C-terminally located cysteine residue to form thioether cyclic compounds analogous to those depicted in Formula I. Other cyclic compounds are prepared in the

same fashion as described elsewhere herein. Also within the scope of this embodiment are amino acid sequence variants of native IgE sequences corresponding to the sequence of this embodiment. Beta strand D variants are selected to enhance binding to FCEH, while the sequence outside of the beta strand D domain need only retain sufficient conformational structure to properly juxtapose the N- and C-termini in substantially the same position as is the case with the native IgE sequence.

[0073] The FCEH-specific polypeptides herein optionally comprise non-IgE polypeptides exactly as described above for the FCEL-specific polypeptides, except that it is not preferred that the FCEH-specific polypeptides comprise cytotoxic functionalities. In addition, conformationally restrained (typically cyclic) polypeptides comprising the FCEH-binding sequence of the beta strand D domain are included within the scope hereof. Such polypeptides are identical to those shown in Formula I above except that the FCEH-binding beta strand D domain replaces the (a3)-(a19) moiety. Exemplary replacement moieties include S420-R428, T421-N430, S420-G433 and R422-R428 (note that sequences such as T421-N430 from U266 that omit a residue from the Kabat sequence can contain a residue at that site or may have a deletion at the same location, in the latter case here the Asn residue would occupy site 429).

[0074] Any one or more of the AB-B domain residues also may be deleted in order to substantially reduce or eliminate FCEL binding. Residue deletion may be preferred for the same reason noted above with respect to the beta strand D domain.

[0075] Examples of candidate AB-B domain substitutional or deletional variants are set forth in the following Table 3. To determine the sequence of each variant, identify the residue for each variant number under each site. For example, the sequence of compound 98 comprises A383 A384 A385, and represents the class of mutations to which mutant 7 belongs.

TABLE 3

AA ¹	HulGE Site			
	350 I	351 R	352 K	353 S
C		55	19	37
M		56	18	38
F		57, 88	8,80	39
Y		4, 75, 83-84,89, 97	7, 73	40
W		58,85	6	41
P		59	1, 74, 78-79	42
G		60, 73	5, 76-77	43
A		61, 98, 102	12, 98-99	44,98,101
V	72	62	13, 97	45
L	73	63	14, 81	46
I	75	64	15,82	47
S		65, 103	16	1-2, 5-36, 55-72, 74, 76-91, 93-95, 97, 99-100, 102, 105
T		66, 104, 105	17	49
D		67, 86	9	50
E		68, 87	89, 94	51
N	79	69	10	52, 79, 84
Q	1-71, 77, 78, 80-95, 97-103, 105	70	11, 103	3, 54, 75, 80, 82-83, 85-89, 103-104
H		71	83, 104	4, 53
K	104	72, 79	2-4, 20-72, 75, 85-88, 91-93, 100-102, 105	48

¹ Amino acid residue substituted into the analogue

TABLE 3 (continued)

AA ¹	HulGE Site			
	350 I	351 R	352 K	353 S
R		1-3, 5-54, 74, 76-78, 80-82, 90-92, 94, 99-101	84	73
Δ^2	96	93, 95, 96	90, 95, 96	92,96

¹Amino acid residue substituted into the analogue²Signifies a deletion

[0076] Insertion of one or more extraneous residues adjacent to a residue within the AB-B domain also falls within the scope of this invention, although substitutions or deletions are preferred. Typically, only one residue will be inserted, although from 2 to 4 or more residues can be inserted adjacent to any one site within the AB-B domain. Smaller numbers of inserted residues will be preferred in order to avoid the introduction of immunogenic sites. This, however, is merely a matter of choice. In general, insertions will be made at a single site, although insertions can be made adjacent to any two or more AB-B domain residues.

[0077] Insertions typically are made between the following residues: S385 and P386, P386 and T387, T387 and I388, and I388 and T389. The inserted residue or residues generally will exhibit charge, bulk or hydrophobicity character which is distinct from that of the flanking residues. For example, candidate insertions can be selected from the following Table 4.

TABLE 4

Insertion	AB-B domain site ¹
Q	1, 2, 3, 4 or 5
D	1, 2, 3, 4 or 5
E	1, 2, 3, 4 or 5
F	1, 2, 3, 4 or 5
W	1, 2, 3, 4 or 5
P	1 or 2
K	2 or 3
R	2 or 3
T	3 or 4
EK	2 or 4
ER	2 or 4
DK	2 or 4
DR	2 or 4
G	1 or 2
A	5
Y	3 or 4
N	1, 2, 3, 4 or 5
H	1, 2, 3, 4 or 5
I	1, 2, 3, 4 or 5

¹I382 - site 1 - R383 - site 2 - K384 - site 3 - S385 - site 4 - P386 - site 5 - T387. Absence of a site indicates no insertion at that site.

[0078] One or more of the AB-B domain residues are substituted or deleted, or additional residues inserted adjacent to such residues. In general, no more than 4 residues or sites are varied and optimally only one will be varied. Variations

herein include combinations of insertions, deletions or substitutions. Excluded from the scope of FCEH specific polypeptides are the linear IgE polypeptide fragments disclosed by Nio et al. (or the naturally occurring sequence variants of such fragments, e.g. alleles and the like), together with any other such fragments disclosed by the prior art.

5 Loop EF Variants

[0079] Loop EF is defined above. Loop EF variants not described in the examples may require screening against both FCEH and FCEL assays since loop EF is involved in both FCEL and FCEH binding. However, this screening will be routine and well within the ordinary skill when following the directions and principles herein. In general, FCEH or FCEL-binding differential polypeptides will comprise substitutions or deletions of (or insertions adjacent to) one or more of residues 446, 447, 448, 449, 450, 452 and 453. It should be noted that sites such as 446 and 447, while shown in the case of Ala substitution to lead to loss of FCEL binding (Example 5), also serve as sites for selecting variants which bind FCEL to a greater degree than native IgE. For the most part, however, sites 446 and 447 are not preferred for introducing variants in which the objective is FCEL binding. For this, one should focus on the region extending from residue 448 to 453, and preferably residues 450, 452 and 453. In general, loop EF variants are employed with variants introduced into loop AB - beta strand B or beta strand D or both.

[0080] R446 typically is substituted by Gly, Ala, Val, Leu, Ile, Ser, His, Lys, Met, Thr, Asp, Pro, Glu, Asn, Gln, Cys, Phe, Tyr or Trp, preferably Ala for FCEH specificity.

[0081] D447 generally is substituted by Gly, Ala, Val, Leu, Ile, Met, Cys, Ser, Thr, Pro, Glu, Asn, Gln, His, Lys, Arg, Phe, Tyr or Trp, preferably Ala for FCEH specificity.

[0082] W448 also generally is not substituted, but if so then Gly, Ala, Val, Leu, Ile, Met, Cys, Ser, Thr, Pro, Glu, Asn, Asp, Gln, His, Lys, Arg, Phe or Tyr are employed.

[0083] I449 likewise generally is not substituted, but if so then Gly, Ala, Val, Leu, Met, Cys, Ser, Thr, Pro, Glu, Asn, Asp, Gln, His, Lys, Arg, Phe, Tyr or Trp are employed.

[0084] E450 typically is substituted with Gly, Ala, Val, Ile, Leu, Met, Cys, Ser, Thr, Pro, Gln, Asn, Asp, His, Lys, Arg, Phe, Tyr or Trp, preferably Ala for FCEH specificity.

[0085] G151 generally is not substituted, but if so then Ala, Val, Leu, Met, Cys, Ser, Thr, Pro, Glu, Asn, Ile, Asp, Gln, His, Lys, Arg, Phe, Tyr or Trp are employed.

[0086] E452 also generally is substituted with Ala, Val, Leu, Met, Cys, Ser, Thr, Pro, Gly, Asn, Ile, Asp, Gln, His, Lys, Arg, Phe, Tyr or Trp.

[0087] T453 typically is substituted with Ala, Val, Leu, Met, Cys, Ser, Pro, Gly, Asn, Glu, Ile, Asp, Gln, His, Lys, Arg, Phe, Tyr, or Trp.

[0088] Exemplary IgE variants are set forth in Table 5. It will be understood that this table may contain variants that bind to both receptors, differentially to one or the other, or to neither receptor.

TABLE 5

AA ¹	HulGE Site				
	446 R	447 D	450 E	452 E	453 T
C	47	46	45	44	43
M	34				
F	33	25			
Y	41				30
W		26	36, 38		36, 38
P			49		
G					
A	13, 17	16	12, 15	12, 14	12
V					31
L					40
I		48			
S					29

¹amino acid residue substituted into the variant

TABLE 5 (continued)

AA ¹	HulGE Site				
	446 R	447 D	450 E	452 E	453 T
T	43				1-3, 5-7, 9, 10, 13-17, 24-26, 28, 33, 34, 37, 39, 44-48, 50, 51
D	39	1, 2, 4-15, 17-23, 31-45, 47, 49-52	5, 8, 11, 18, 23, 27, 32, 33, 35, 40, 42, 52	1, 29, 30, 34, 50	42
E	9, 20	24, 29, 30	1-5, 7, 9, 10, 13, 14, 16, 17, 24-28, 30, 31, 34, 37, 39, 43, 44, 46, 47, 48, 51	3, 4, 6, 7, 9, 10, 13, 15-17, 24-26, 28, 31-33, 37, 39, 43, 45-49, 52	8, 11, 18-23, 27, 35
N	19, 22, 40	3	50	51	
Q	10, 11, 23, 35, 36, 42			2	52
H	21, 30	27		36	
K	18, 28, 29, 52	28		8, 11, 18-23, 27, 35, 40, 42	32
R	1-8, 12, 14-16, 24-27, 31, 32, 38, 44-46, 48-51	7	6	5	4
Δ^2	37			38	

¹amino acid residue substituted into the variant²signifies a deletionVariant Anti-hulGE antibodies

[0089] Variant anti-hulGE antibodies were produced by first obtaining a group of murine monoclonal antibodies which were capable of binding to FCEL but not to FCEH. 8 such murine monoclonal antibodies, designated MAE10, MAE11, MAE12, MAE13, MAE14, MAE15, MAE16 and MAE17, were obtained by conventional methods involving immunizing mice with human IgE or a polypeptide consisting of residues 315-547 of hulGE and screening for anti-IgE activity.

[0090] MAE11/15 and MAE13 recognize different epitopes. It appears that the MAE13 epitope is located three-dimensionally adjacent to a key component of the FCEH binding site of IgE (but does not directly occupy that site) since a slight amount of histamine release will occur at high concentrations of MAE 13 suggesting that some limited antibody mediated crosslinking of FCEH occurs with MAE 13. MAE17 was most effective in suppressing B-cell IgE synthesis despite the fact that MAE11 and MAE13 exhibited greater IgE affinity. This may be attributed to its ability to mediate complement fixation (it possessed an IgG2a isotope, thus containing an Fc capable of eliciting effector function).

[0091] MAE11 and MAE15 are believed to recognize the same IgE epitope. Each antibody shared certain unusual features in its amino acid sequence. For example, CDR1 of the light chain of each contained 3 aspartic acid residues. CDR3 of the heavy chains of MAE11 and MAE15 contained 3 histidine residues (and contained two arginine residues, respectively).

[0092] Antibodies such as the foregoing having desired IgE binding characteristics may be further modified. Such modifications fall into two general classes. In the first class the antibodies are modified so that they are monovalent for IgE. This means that only one "arm" of the antibody, i.e., one light-heavy chain fork of the antibody, shall be capable of binding IgE. The remaining Fv "arm" of the antibody (or arms in the case of IgM) is specific for a second (non-IgE) antigen, is not capable of binding any antigen, or is deleted entirely. Thus, the term IgE monovalent covers polyvalent antibodies that are monovalent for IgE. The best results may be obtained with the second alternative, since this would preserve the structure of the antibody most faithfully and would likely confer the longest circulating half-life on the antibody. IgE-monovalent antibodies specific for FCEL bound IgE optimally will comprise sufficient fc domains of the

heavy chains to be capable of complement binding and Ig effector functions.

[0093] The second antigen recognized by one embodiment of IgE monovalent antibody is one which, when indirectly cross-linked to FCEL by the antibody herein, will not produce any toxic or deleterious response, i.e. the second antigen is not FCEH, and generally is one which is not found in the animal to be treated (in order to avoid undesired absorption of the antibody onto tissues or proteins within the body). Thus, the second antigen ordinarily will not (but may be) FCEL. However, in some circumstances the second antigen will be a protein present in the patient to be treated, e.g. where such proteins are to serve as carriers or depot releases for the therapeutic antibodies herein.

[0094] Such IgE-monovalent antibodies are made by methods known per se. For example, DNA encoding the anti-IgE Fv heavy and light chains is ligated to DNA encoding the Fc of a human recipient antibody. In addition, DNA is provided that encodes heavy and light chains for an antibody capable of binding second antigen or an unidentified antigen, or that encodes heavy and light chain having sufficient residues deleted from the CDRs that non-IgE antigen binding no longer can occur. A conventional recombinant host is transformed with all four DNAs and the products recovered. Assuming random chain assortment, a subpopulation of antibody products will contain one arm with anti-IgE heavy and light chain and at least another arm having specificity for second antigen or no antigen. The desired subpopulation then is purified by conventional methods, e.g., immunoaffinity absorption or by molecular sieving. These antibodies also can be made by reduction of the starting antibodies followed by oxidative chain recombination, as has heretofore been employed in the preparation of monovalent antibodies (see for example Glennie *et al.*, *Nature* 295: 712 [1982]).

[0095] In addition to IgE-monovalency, in other embodiments the antibodies are modified so that they contain a maximum proportion of human sequence (commensurate with retention of required or desired activity), i.e., they are converted to chimeras or are humanized. In both instances the functional effect is to place the anti-IgE binding capability of the murine or other donor antibody into a human background to make it as non-immunogenic as possible. General methods are known for making chimeras and for humanizing antibodies (as noted above). A minimal amount of non-human antibody sequence is substituted into the recipient human antibody. Typically, the non-human residues are substituted into the V_H, V_L, V_H-V_L interface or framework of the recipient human antibody. Generally, the Kabat CDR's of the humanized antibodies are about 80% and more typically about 90% homologous with the non-human donor CDR's. The V_H-V_L interface and framework residues of the humanized antibody, on the other hand, are about 80%, ordinarily 90% and preferably about 95% homologous with the recipient human antibody. Homology is determined by maximal alignment of identical residues. The resulting antibody is (a) less immunogenic in humans than a murine antibody and (b) capable of binding to FCEL-bound hulgE but substantially incapable of binding to FCEH-bound hulgE. Such antibodies typically comprise a human antibody which is substituted by an amino acid residue from a complementarity determining region (CDR), VL-VH interface or a framework region of a non-human anti-IgE antibody which is capable of binding. One or more, and preferably all, of the nonhuman CDR's L1, L2, L3, H1, H2 or H3 are substituted into the human antibody recipient.

[0096] The characteristics possessed by the MAE11 antibody were preferred for therapeutic use. Since MAE11 bound to soluble IgE, bound to MlgE bearing B cells, blocked IgE binding to the low and high affinity IgE receptor, inhibited *in vitro* IgE production and failed to bond to IgE coated basophils, it was chosen as the donor antibody for humanization. The recipient antibody was Kabat human kappa (light) subgroup I and human subgroup III heavy chain, although it will be understood that any other human antibody can be suitably employed. Surprisingly, optimal results were not obtained by simply substituting the murine CDRs in place of the CDRs in a recipient human antibody (Fig. 3; Table 8 *infra*). Instead, it was necessary to restore donor framework hydrophobic residues such as V_H 78, 48, 49, 63, 67, 69; 82 or 82c, or V_L 13, 19, 58, 78 or 104, in order to achieve a degree of inhibition of IgE binding similar to that of the donor antibody. While these residues function to establish the conformation of CDRs, they generally are not exposed to the exterior of the antibody so use of the murine residues should not exert a significant impact on immunogenicity. Other non-CDR residues exerting an effect on binding included V_H60, 61, 37, 24, and V_H50, 52, 58 and 95 (non-CDR by Chothia), and V_L4, V_L33 (non-CDR by Chothia) and V_L53 (non-CDR by Chothia). The human framework hydrophobic residues generally are substituted with other hydrophobic residues (especially those from the donor antibody) such as valine, isoleucine, leucine, phenylalanine or methionine. The remaining non-CDR residues are substituted with any other amino acid residue, but again preferably the murine residue found at the analogous site.

[0097] In general, the character of the anti-IgE antibody is improved by substituting, deleting or inserting a residue at or adjacent to V_L sites 29a, 29c, 30, 33, 55, 57, 58, 78, 93, 94, or 104 and/or V_H residues 24, 37, 48, 49, 54, 57, 60, 61, 63, 65, 67, 69, 78, 82, 82c, 97, 100a or 100c.

[0098] Position V_H-78 is most preferably substituted with phenylalanine. However, it also is substituted with leucine, valine, isoleucine, methionine, alanine or any other residue which results in an improvement in the characteristics of the antibody (see *infra*).

[0099] Position V_H-60 is most preferably substituted with asparagine, although substitution with glutamine, histidine, lysine, arginine or any other residue which improves the characteristics of the antibody shall fall within the scope of this invention.

[0100] Position V_H-61 is most preferably substituted with proline, although glycine, alanine, valine, leucine, isoleucine or any other residue which results in an improvement in the characteristics of the antibody also is suitable.

[0101] CDR residues were imported from the donor MaE11. These included four inserts in V_L1, 30a-30d, as well as 91-94 (V_L1), V_H1 27-29, 29a, 31, 33 and 34, V_H2 53-55, and V_H3 97-101. V_L 29a, 29c or 30, as well as V_H97, 100a or 100c, are important in conferring on the CDR ability to bind IgE.

[0102] V_H positions 97, 100a and 100c in humae11 (humanized Mae11) are all histidine, and 2 are arginine in MaE15. These residues are important in IgE binding. One, two or three of these are modified by substitution with basic residues, particularly lysine or arginine, but also with alanine, glycine, valine, isoleucine, serine, threonine, aspartic acid, glutamic acid, asparagine, glutamine, methionine, phenylalanine, tyrosine, tryptophan or proline.

[0103] V_L positions 29a, 29c and 30 of humae11 also are important for IgE binding. In humae11 each of these positions are occupied by the acidic residue, aspartic acid. They are substituted in other embodiments by glutamic acid, but also may be substituted with alanine, glycine, valine, isoleucine, serine, threonine, asparagine, glutamine, methionine, phenylalanine, tyrosine, tryptophan or proline. It is within the scope of this invention to reverse the charges on positions V_L 29a, 29c and 30 with those on V_H 97, 100a and 100c, e.g. by employing aspartic acid residues in the three V_H sites (2 in the case of humanized MaE15) and histidine in the three V_L sites.

[0104] Residues also may be inserted adjacent to V_H positions 97, 100a, 100c, 61 or 61, or V_L residues at positions 29a, 29c, 30 or 78. Inserted residues generally will be of like kind, e.g. an acid residue would be inserted adjacent to V_L-29a, 29c or 30, while a basic residue is inserted adjacent to V_H-97, 100 or 100c. The residues at these sites also may be deleted.

[0105] Humanized IgE-monovalent antibodies also are included within the scope of this invention. In this instance humanization extends to the anti-IgE arm as well, if necessary, to the remaining arm(s). Non-IgE binding arms of course can originate from human antibodies and in such case will not require humanization.

[0106] The foregoing variations are made by introducing mutations into the DNA encoding the precursor form of the antibody and expressing the DNA in recombinant cell culture or the like. This is accomplished by conventional methods of site directed mutagenesis. The variants then are screened for the desired character in assays conventional *per se*. In the case of anti-huIgE, desired character includes increasing the antibody affinity for huIgE, increasing its capacity and specificity for FCEL bound IgE, increasing the concentration of antibody required to stimulate histamine release from mast cells or basophils, reducing immunogenicity in humans, and other improvements apparent to the ordinary artisan. Optimizing these characteristics frequently will require balancing one improvement against another and therefore is a matter of judgment and is dependent upon the performance parameters dictated by the use intended for the antibody.

[0107] It is preferable to use a human IgG1 (or other complement fixing antibody) as the recipient immunoglobulin for humanization, although hu IgG2, IgG3, IgG4, IgE, IgM, IgD or IgA also can be used as recipient. Preferably the recipient is a complement fixing IgG antibody or an IgG antibody capable of participating in ADCC.

Therapeutic, Diagnostic and Preparatory Uses

[0108] The anti-IgE antibodies herein are useful in identifying IgE amino acid sequence variants in which the FCEL or FCEH-binding domains have been modified. Candidate FCEL or FCEH-specific polypeptides are incubated with these antibodies, and analogues to which these antibodies fail to bind are selected for further evaluation, e.g., determination, respectively of their FCEH and FCEL receptor binding characteristics. Any antibody, whether of murine, human, or another animal species in origin, or a variant thereof such as the humanized immunoglobulins described above, which has the epitopic specificity of any of antibodies MAE10 - MAE17 (especially MAE11/15, MAE13 or MAE17) will be equally acceptable. Such antibodies are easily identified by immunizing a suitable animal or using an *in vitro* Fv selection system, e.g. phagemid, with IgE of the appropriate animal origin and screening the animals or products for antibodies having the ability to compete for IgE with MAE11/15, 13, 17 or other antibodies which map to substantially the same epitopic site(s) as those described herein. As noted, the antibodies desirably are monovalent for FCEL-bound IgE when employed therapeutically. They may be bivalent and/or bispecific when used to purify IgE from plasma, serum or recombinant cell culture.

[0109] The FCEH and FCEL-specific, differential binding polypeptides are useful for diagnostics and therapeutics. In *in vitro* diagnostic assays they are employed as specific binding reagents in assays for FCERI or FCERII, respectively. The polypeptides of this invention are labelled with a detectable substance such as an enzyme, fluorescent or chemiluminescent group, radioisotope or a specific binding moiety that binds to a detectable substance (such as an enzyme). A typical specific binding moiety is an immunoglobulin variable domain which is capable of binding to the detectable substance. FCEL and FCEH specific polypeptides comprising immunoglobulin variable domains are described in more detail above.

[0110] Assay systems that employ the FCEL or FCEH specific polypeptides of this invention are analogous to the sandwich-type systems heretofore generally used in the immunoassay field. Here, the specific polypeptide is employed

in the same fashion as labelled antibodies directed against antigen (the FCEH or FCEL receptor) or as an absorption agent insolubilized on a matrix for the isolation of receptor from test sample. Redox, proteolytic, esterolytic or other conventional enzyme labels are conjugated to the polypeptides of this invention for use in conventional assay systems.

[0111] The differential binding polypeptides of this invention also are useful for the isolation of FCEL or FCEH from cell culture in preparing FCEL or FCEH for therapeutic or research purposes. The polypeptide is covalently bonded or noncovalently adsorbed to a matrix such as an ion exchange resin, an immunoaffinity column (containing an antibody capable of binding a polypeptide fused to the FCEH or FCEL-specific polypeptide), an immobilized antigen (where the FCEH or FCEL-specific polypeptide comprises an immunoglobulin variable region capable of binding to the antigen) or a cyanogen bromide activated polysaccharide. The immobilized FCEH or FCEL-specific polypeptide then is contacted with the receptor preparation under conditions such that the receptor is bound to the FCEH or FCEL-specific polypeptide. The receptor then is eluted by changing the pH or ionic conditions and separating the polypeptide preparation from the receptor.

[0112] The differential binding polypeptides herein are useful in preparing antibodies specific to the FCEH or FCEL-binding domain of IgE. For example, antibodies capable of binding specifically to the FCEH or FCEL-binding domains of IgE are selected by first immunizing a subject with IgE. Monoclonal antibodies then are selected in the ordinary way for native IgE binding, and the monoclonal antibodies then screened to identify those that bind to a FCEH or FCEL-specific polypeptide of this invention. Preferably the FCEH or FCEL-specific polypeptide will be identical in sequence to the corresponding sequence of the IgE used as immunogen except, of course, for the minimal mutations need to confer FCEH or FCEL differential binding specificity. For example, the IgE monoclonal antibodies can be selected for their inability to bind to mutation 6. If they are unable to bind to mutation 6 one can conclude that they bind to the FCEH-binding site and are therefore promising for use in diagnostic or therapeutic procedures that depend upon an antibody that fails to bind to FCEH-bound IgE but which binds to FCEL-bound IgE. Confirmation is obtained by determining that the antibody selected in fact binds to IgE bound to FCEL. Since the selected antibody is highly specific for the key site(s) involved in receptor binding it is then possible to reduce the size of the antibody; the bulk of the antibody is not needed for steric hinderance of the IgE-receptor interaction. Thus, it becomes feasible in allergy therapy to use anti-IgE monovalent antibodies or other anti-IgE fragments such as Fab, Fab' and the like.

[0113] Similarly, the FCEL or FCEH-specific polypeptides are useful as immunogens for raising antibodies capable of cross-reacting with native IgE only at epitopic sites outside of the domains varied in creating the FCEH or FCEL-specific polypeptides. For example, mutations 6 and 7 are useful for raising antibodies specific for IgE epitopes except for the mutated AB-B or beta strand B domains as the case may be.

[0114] The FCEH and FCEL-specific polypeptides and anti-IgE antibodies (especially those with reduced immunogenicity) are useful in therapies for the treatment or prophylaxis of allergies, although the FCEH specific polypeptide subgroup which bears cytotoxic functionalities is not considered suitable for therapy since it could lead to degranulation of mast cells and basophils. Otherwise, the polypeptides typically are administered to a patient who is known to be sensitized to an allergen, preferably prior to an acute allergic response. The dosages and administration route will depend upon the accessory functionalities accompanying the polypeptides (e.g. cytotoxic agents, immunoglobulin effector functions, etc.), the condition of the patient (including the population of B cells or mast cells and basophils), the half-life of the polypeptide, the affinity of the polypeptide for its receptor and other parameters known to the clinician. As a general guide in the case of FCEH-specific polypeptide, one will determine from blood tests the amount of target cells circulating in the patient and determine the amount of polypeptide to displace or effectively compete with endogenous IgE taking into account the population of FCEH receptors as well as the half life and affinity of the polypeptide for FCEH. An excess of polypeptide calculated to be necessary to substantially displace native FCEH-bound IgE over a reasonable therapeutic interval will then be administered. Similar analysis used to determine the dosage of anti-IgE antibody or FCEL polypeptide.

[0115] Therapeutic polypeptides are administered by intravenous intrapulmonary, intraperitoneal subcutaneous or other suitable routes. Preferably the polypeptides are administered s.c. or i.v. over a period of about from 1 to 14 days as required. In the case of FCEL-specific polypeptide or anti- FCEL- bound IgE one would determine the amount needed to inhibit, suppress or kill a substantial portion of the IgE-secreting B cell population. Inhibition or suppression of the B cell population includes either or both of reductions in IgE secretion and attenuation of the total number of IgE secreting B cells. Candidate doses are readily determined by the use of *in vitro* cell cultures or animal models.

[0116] Therapy of allergic disorders with anti- FCEL bound IgE and FCEL or FCEH polypeptides optionally is accomplished with other known therapies for allergies. These include administration of gamma interferon, allergen desensitization, reduction in exposure to allergen, treatment with antihistamines and the like.

Preparation of FCEH- and FCEL-Specific Polypeptides

[0117] The FCEH- or FCEL-specific polypeptides of this invention are made in conventional fashion, i.e., modifications of amino acid sequence are accomplished by commonly available DNA mutagenesis methods such as PCR

amplification using primers bearing the mutants, or by M13 mutagenesis, followed by expression of the mutated DNA in recombinant host cells. The polypeptides also can be made by Merrifield or other *in vitro* methods of synthesis if they are sufficiently small (generally, under about 100 residues). However, the polypeptides preferably are made by recombinant methods. Selection of recombinant host cells, vectors, culture conditions and other parameters are not believed to be critical. In general, hosts, vectors and methods heretofore used in the recombinant expression of immunoglobulins (generally, IgGs) are also useful for the preparation of the polypeptide sequences of this invention. Preferably, mammalian cells such as myelomas, CHO, Cos, 293s and the like are employed as hosts, and the vectors are constructed for secretory expression of the polypeptide. Recombinant expression systems facilitate the preparation of functional immunoglobulin variants containing FCEL- or FCEH-specific sequences since the host cells can be transformed with DNA encoding one heavy chain containing the FCEL- or FCEH-specific sequences and one light chain, each of which contains a variable domain for binding a first antigen, and an immunoglobulin that binds antigen and FCEL or FCEH recovered. Similarly, the same process is used with DNA encoding in addition another heavy chain containing the FCEL- or FCEH-specific domain and another light chain, each of which contain a variable domain for binding a second antigen, and a bivalent immunoglobulin recovered. Properly assembled immunoglobulin analogues are recovered by affinity chromatography on a matrix containing the two antigen(s).

[0118] The polypeptides of this invention are recovered from lysed recombinant cell culture or (when secreted) the culture supernatant. Substantial purification is achieved by passing cell free extracts which contain the polypeptides over an immobilized FCEL or FCEH receptor affinity matrix. Other methods heretofore used to purify IgE or other appropriate immunoglobulins are equally acceptable here, including immunoaffinity and (when appropriate) absorption on immobilized antigen.

[0119] Polypeptides of this invention which contain short sequences preferably are prepared using solid-phase synthesis, e.g. the method of Merrifield, *J. Am. Chem. Soc.*, **85**:2149 (1963). However, other equivalent chemical syntheses known in the art are acceptable. The recombinant or *in vitro* synthesized polypeptides then are cross-linked to matrices (for use in diagnostic or preparatory procedures) or are placed into conformationally restrained structures. Known cyclizing procedures such as those described in PCT 90/01331 or Lys/Asp cyclization using N α -Boc-amino acids on solid-phase support with Fmoc/9-fluorenylmethyl (Ofm) side-chain protection for Lys/Asp, followed by piperidine treatment and cyclization, are useful. Methods which depend upon cross-linking or cyclization through residue side chains may require that an extraneous residue be inserted at the C and/or N terminus of the AB-B or beta stand D domains, as the case may be, to provide a suitable cyclizing or cross-linking site.

[0120] Glu and Lys side chains also have been crosslinked in preparing cyclic or bicyclic peptides: the peptide is synthesized by solid phase chemistry on a p-methylbenzhydrylamine resin, the peptide is cleaved from the resin and deprotected. The cyclic peptide is formed using diphenylphosphorylazide in diluted methylformamide. For an alternative procedure, see Schiller *et al.*, *Peptide Protein Res.* **25**:171-77 (1985). See also U.S. Patent 4,547,489.

[0121] Disulfide crosslinked or cyclized peptides are generated by conventional methods. The method of Pelton *et al.*, *J. Med Chem.*, **29**:2370-2375 (1986) is suitable. Also useful are thiomethylene bridges (*Tetrahedron Letters* **25**: 2067-2068 (1984). See also Cody *et al.*, *J. Med Chem.*, **28**:583(1985). The C390 residue found in the C-terminal sequence of the AB-B domain is useful in cross-linking or cyclizing this domain.

[0122] Typically, extraneous residues which are to participate in cyclization or cross-linking are inserted at the N- and C-termini of the chosen AB-B or beta strand D sequence as part of the synthesis of the polypeptide precursor to be employed in the procedure. The desired cyclic or cross-linked peptides are purified by gel filtration followed by reversed-phase high pressure liquid chromatography or other conventional procedures. The peptides are sterilized by 0.2 μ m filtration and formulated into conventional pharmacologically acceptable vehicles.

[0123] The compounds described in this invention may be the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of this invention. Examples of such salts include ammonium, metal salts like sodium, potassium, calcium and magnesium; salts with organic bases like dicyclohexylamine-N-methyl-D-glucamine and the like; and salts with amino acids such as arginine or lysine. Salts with inorganic and organic acids may be like prepared, for example, using hydrochloric, hydrobromic, sulfuric, phosphoric, trifluoroacetic, methanesulfonic, maleic, fumaric and the like. Non-toxic and physiologically compatible salts are particularly useful although other less desirable salts may have use in the processes of isolation and purification.

[0124] A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. For example, reaction of the free acid or free base form of a compound of Formula I with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble; or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free acid or base form of the product may be passed over an ion exchange resin to form the desired salt, or one salt form of the product may be converted to another using the same general process.

[0125] Additional pharmaceutical methods may be employed to control the duration of action of the polypeptides of this invention. Controlled release preparations are achieved through the use of polymers which complex with or absorb the subject polypeptides. Controlled delivery is achieved by formulating the polypeptides into appropriate macromo-

lecular articles (for example, those prepared from polyesters, polyamino acids, polyvinyl, polypyrrolidone, ethylene vinylacetate, methylcellulose, carboxymethylcellulose, or polyamine sulfate).

[0126] Alternatively, instead of entrapping the polypeptides in polymeric matrices, it is possible to entrap these materials in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization. Hydroxymethylcellulose or gelatin microcapsules and poly(methylmethacrylate) microcapsules, respectively, are useful, as are in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nanoparticles and nanocapsules). See Remington's Pharmaceutical Sciences (1980).

EXAMPLE 1

Preparation of monoclonal antibodies to IgE

[0127] Eight monoclonal antibodies with the ability to block the binding of IgE to the FCEH were used. These monoclonal antibodies, referred to as MAE10 - MAE17, were made in the following manner. Purified human IgE was prepared from supernatants of U266B1 cells (ATCC TIB 196) using affinity chromatography on a previously isolated anti-IgE antibody (Genentech MAE1, although other anti-huIgE antibodies are equally useful). For MAE12, five BALB/c female mice, age six weeks, were immunized in their foot pads with 10 µg of the purified IgE in Ribi's adjuvant. Subsequent injections were done in the same manner one and three weeks after the initial immunizations. Three days after the final injection, the inguinal and popliteal lymph nodes were removed and pooled, and a single cell suspension was made by passing the tissue through steel gauze. For MAE14, MAE15, and MAE13 the immunizations were done in a similar manner except that for MAE13 30 µg of IgE per injection were used and IgE 315-547 was used as a prefusion boost; for MAE14 and MAE15 five injections of 50 µg each were used; and the IgE immunogen for MAE17 was IgE 315-547. For MAE10 and MAE11, injections were given subcutaneously in two doses of 100 µg and a final booster of 50 µg, and spleen cells were used for the fusions. The cells were fused at a 4:1 ratio with mouse myeloma P3X63-Ag8.653 (ATCC CRL 1580) in high glucose (DMEM) containing 50% w/v polyethylene glycol 4000.

[0128] Fused cells were plated at a density of 2×10^5 per well in 96 well tissue culture plates. After 24 hours HAT selective medium (hypoxanthine/aminopterin/thymidine, Sigma Chemical Company, # H0262) was added. Of 1440 wells plated, 365 contained growing cells after HAT selection.

[0129] Fifteen days after the fusion, supernatants were tested for the presence of antibodies specific for human IgE using an enzyme-linked immunosorbent assay (ELISA). The ELISA was performed as follows, with all incubations done at room temperature. Test plates (Nunc Immunoplate) were coated for 2 hours with rat anti-mouse IgG (Boehringer Mannheim, # 605-500) at 1 µg/ml in 50 mM sodium carbonate buffer, Ph 9.6, then blocked with 0.5% bovine serum albumin in phosphate buffered saline (PBS) for 30 minutes, then washed four times with PBS containing 0.05% Tween 20 (PBST). Test supernatants were added and incubated two hours with shaking, then washed four times with PBST. Human IgE (purified from U266 cells as described above) was added at 0.5 µg/ml and incubated for one hour with shaking, then washed four times in PBST. Horseradish peroxidase conjugated goat anti-human IgE (Kirkegaard & Perry Labs, # 14-10-04, 0.5 mg/ml) was added at a 1:2500 dilution and incubated for one hour, then washed four times with PBST. The plates were developed by adding 100 µl/well of a solution containing 10 mg. of o-phenylenediamine dihydrochloride (Sigma Chemical Company # P8287) and 10 µl of a 30% hydrogen peroxide solution in 25 ml of phosphate citrate buffer Ph 5.0, and incubating for 15 minutes. The reaction was stopped by adding 100 µl/well of 2.5 M sulfuric acid. Data was obtained by reading the plates in an automated ELISA plate reader at an absorbance of 490 nm. For MAE12, 365 supernatants were tested and 100 were specific for human IgE. Similar frequencies of IgE specificity were obtained when screening for the other antibodies. All of the monoclonal antibodies described herein were of the IgG1 isotype except for MAE17, which was IgG2b, and MAE14, which was IgG2a.

[0130] Each of the IgE specific antibodies was further tested in cell-based and plate assays to select for antibodies which bound to IgE in such a way as to inhibit IgE binding to FCEH and which are not capable of binding to FCEH-bound IgE. The results of these assays are set forth in Table 5 and Table 5a below.

TABLE 5
SUMMARY OF MURINE Anti-Eu IgE mAb CHARACTERISTICS

mAb	Immunogen	Schedule/ Dose (μ g)	B-cell source	Isotype	† Binding FCRH-bound IgE ¹	PBL Histamine Release ¹ (BC50)	Amount blocking FCRH ¹ (BC50)
MaB 1	PS IgE	3x50	Lymph Node	IgG1	.05 μ g/ml	1 μ g/ml	0.3 μ g
MaB 10	U266 IgE	2x100, 1x50	Spleen	IgG1	No binding at 10 μ g/ml	>100 μ g/ml	2.5 μ g
MaB 11	U266 IgE	2x100, 1x50	Spleen	IgG1	No binding at 10 μ g/ml	>100 μ g/ml	0.6 μ g
MaB 12	U266 IgE	3x30	Lymph Node	IgG1	No binding at 10 μ g/ml	>100 μ g/ml	0.8 μ g
MaB 13	U266 IgE	3x30	Lymph Node	IgG1	No binding at 10 μ g/ml	>10 μ g/ml	0.6 μ g
MaB 14	U266 IgE	5x50	Lymph Node	IgG2a	No binding at 10 μ g/ml	>100 μ g/ml	2.5 μ g
MaB 15	U266 IgE	5x50	Lymph Node	IgG1	No binding at 10 μ g/ml	>100 μ g/ml	0.6 μ g
MaB 16	rHIgE aa 315- 547	5x1	Lymph Node	IgG1	No binding at 10 μ g/ml	>100 μ g/ml	0.7 μ g
MaB 17	rHIgE aa 315- 547	5x1	Lymph Node	IgG2b	No binding at 10 μ g/ml	>100 μ g/ml	>5.0 μ g

Table 5a. Summary of murine Anti-Hu IgE mAb (continued)

mAb	† Binding to Membrane IgE on U266BL ₁ (BC50) ⁴	† Binding of IgE on FcεRI (CD23) IM9 (BC50) ⁵	Blocks 1μg IgE binding to FcεRI II (BC 50) ⁶	Inhibition of in-vitro IgE synthesis ⁷	Affinity Constant for IgE ⁸ (Kd)
MaE 1	0.4 μg/ml	.05 μg/ml	>100 μg	(-)	5.4x10 ⁻⁴
MaE 10	0.5 μg/ml	No binding at 10 μg/ml	2.5 μg	(-)	7x10 ⁻⁹
MaE 11	0.15 μg/ml	No binding at 10 μg/ml	0.6 μg	(+)	3x10 ⁻⁴
MaE12	>10 μg/ml	1 μg/ml	5.0 μg	(-)	4x10 ⁻⁷
MaE 13	1 μg/ml	No binding at 10 μg/ml	0.7 μg	(++)	5x10 ⁻⁴
MaE 14	6 μg/ml	No binding at 10 μg/ml	2.5 μg	(±)	1.4x10 ⁻⁴
MaE 15	6 μg/ml	No binding at 10 μg/ml	0.6 μg	(±)	7x10 ⁻⁴
MaE 16	10 μg/ml	<.05 μg/ml	5 μg	(+)	ND
MaE 17	10 μg/ml	No binding at 10 μg/ml	5 μg	(++)	ND

1. FACS based assays for analysis of murine anti-human IgE monoclonals. Screen of murine anti-human IgE monoclonal binding to IgE on CHO 3D10 (FcεRI alpha +)

[0131]

a. CHO 3D10 cells (FcεRI alpha chain stable transfectant; Hakimi *et al.*, *J. Biol. Chem.* **265**:22079) at 5 x 10⁵ cells per sample are incubated with U266 IgE standard (lot no. 13068-46) at 10 μg/ml in 100 μl FACS buffer (0.1% BSA 10mM sodium azide in PBS pH 7.4) for 30 minutes at 4°C followed by one wash with FACS buffer. The amount of

IgE binding is determined by incubating an aliquot of IgE loaded cells with a polyclonal FITC conjugated rabbit anti-human IgG (Accurate Chem. Co. AXL-475F, lot no 16) at 50 µg/ml for 30 minutes at 4°C followed by three washes with FACS buffer.

b. IgE loaded cells are incubated with 100 µl of murine anti-human IgE hybridoma supernatant (murine IgG concentration ranging from 1 to 20 µg/ml) for 30 min. at 4°C followed by one wash with FACS buffer. A Genentech monoclonal anti-human IgE (MAE1) at 10µg/ml is used as a positive control for binding. Genentech monoclonal (MAD 6P) which does not recognize IgE is used at 10µg/ml as a negative control.

c. Monoclonal binding to human IgE on CHO cells is detected by incubating cells with 20 µg/ml FITC-conjugated, affinity purified F(ab) 2 Goat anti-mouse IgG (Organon Teknica cat. no. 10711-0081) for 30 minutes at 4°C followed by three washes with FACS buffer. Cells are added to 400µl buffer contain 2 µg/ml propidium iodide (Sigma cat no. P4170) to stain dead cells.

d. Cells are analyzed on a Becton Dickinson FACSCAN flow cytometer. Forward light scatter and 90 degree side scatter gates are set to analyze a homogeneous population of cells. Dead cells which stain with propidium iodide are excluded from analysis. Hybridoma supernatants which do not bind IgE on CHO 3D10 cells were considered candidates for further screening.

2. Histamine release from peripheral blood basophils;

[0132] Heparinized blood was obtained from normal donors and diluted 1:4 in a modified Tyrodes buffer (25mM tris, 150mM NaCl, 10mM CaCl₂, MgCl₂, 0.3 mg/ml HSA, pH 7.35) then incubated with 1nM human IgE (ND) at 4°C for 60 minutes. Cells were then added to Tyrodes buffer containing either the murine monoclonal anti-IgE Abs (10 mg/ml) or a polyclonal anti-human antiserum as the positive control, and incubated at 37°C for 30 minutes. Cells were pelleted, histamine in supernatants was acetylated and histamine content was determined using an RIA kit (AMAC, Inc. Westbrook, Main). Total histamine was determined from cells subjected to several rounds of freezed thawing. Percent histamine release was calculated as nM histamine content in supernatant - nM histamine spontaneously released divided by nM total histamine in the sample.

3. Blocking of Fitc conjugated IgE binding to FcERI alpha chain.

[0133] The effect of the antibodies on IgE binding was studied by preincubating Fitc labelled IgE with the various Mae antibodies at 37° C for 30 minutes in PBS containing 0.1% BSA and 10mM Sodium Azide pH 7.4, then incubating the complex with 5×10^5 3D10 cells at 4°C for 30 minutes. The cells were then washed three times and mean channel fluorescence at 475 nM was measured. A murine anti-human IgE mAb (Mae1) which does not block IgE binding to the FcERI alpha chain was used as a control.

4. Analysis of murine anti-human IgE binding to membrane IgE positive B cell U266

[0134]

a. U266 B1 cells (membrane IgE +) are cultured in base medium supplemented with 15% head inactivated fetal calf serum (Hyclone cat no. A-1111-L), penicillin, streptomycin (100 units/ml) and L-glutamine (2mM).

b. Cells (5×10^5 /aliquot) are incubated in 100µl FACS buffer containing murine anti-Human IgE monoclonals at 10, 5, 1, 0.5, and 0.1µg/ml for 30 minutes on ice in 96 well round bottom microtiter plates followed by two washes with FACS buffer. The Genentech monoclonal MAE1 is used as a positive control.

c. Cells are incubated in 100µl FACS buffer containing 50µg/ml (1:20 stock) FITC conjugated F(ab') 2 affinity purified goat anti-mouse IgG (Organon Teknika Cat. no. 1711-0084) for 30 minutes on ice followed by three washes with FACS buffer. Cells are added to 400 µl FACS buffer containing propidium iodide at 2 µg/ml to stain dead cells.

5. FACS based binding assays to FcERII(CD23+) B cell IM9

[0135]

a. FACS analysis of IgE binding to FcERII(CD23) (+) B cell line IM9. The IM9 human B cell myeloma ATCC CCL 159. (Ann. N.Y. Acad. Sci., 190:221-234 [1972]) was maintained in GIF base medium with 10% heat inactivated fetal bovine serum, penicillin, streptomycin (100 units/ml) and L-glutamine (2mM).

b. Cells (5×10^5 aliquot) were incubated in 100µl of FACS buffer containing U266 IgE standard at 2µg/ml for 30 minutes at 4°C in 96 well microtiter plates followed by 2 washes with FACS buffer. As a control, cells were incubated in buffer alone or buffer containing 2µg/ml human IgG1 (Behring Diagnostics Cat. no. 400112, lot no. 801024).

c. Cells were then incubated with murine anti-human IgE monoclonals at 0.1 to 10µg/ml for 30 minutes on ice. Genentech monoclonal MAE1 was used as a positive control.

d. Cells were incubated in 100µl FACS buffer containing FITC conjugated F(ab¹)₂ goat anti-mouse IgG at 50 µg/ml (Organon Teknika Ca #1711-0084) for 30 minutes at 4°C followed by 3 washes with FACS buffer.

e. Cells were added to 400µl buffer containing propidium iodide at 2µg/ml to stain dead cells.

f. Cells were analyzed on a Becton Dickinson FACSCAN flow cytometer. Forward light scatter and 90 degree side scatter gates were set to analyze a homogeneous population of cells and dead cells which stained with propidium iodide were excluded from analysis. FITC positive cells (IgE binding) were analyzed relative to cells stained with FITC rabbit anti-Human IgE alone.

g. As a positive control to determine the level of CD 23 on the surface of IM9 cells in each experiment, an aliquot of cells was stained with Becton Dickinson murine monoclonal Leu 20 (anti-CD23) at 10µg/ml for 30 minutes at 4°C followed by 2 washes. The cells were then incubated with FITC conjugated f(ab')₂ affinity purified goat anti-murine IgG at 50µg/ml.

6. Antibody blocking of Fite conjugated IgE binding to the low affinity IgE receptor.

[0136] The binding of 40 nM FITC labelled IgE to the low affinity IgE receptor (CD23) expressed on the B lymphoblast cell IM-9 was analyzed by flow cytometry on a FACSCAN flow cytometer. The effect of the antibodies on Fite IgE binding was studied by preincubating Fite IgE with the murine anti-human antibodies at 0.1 to 10µg/ml. chimera at 37°C for 30 minutes in PBS containing 0.1% BSA and 10mM Sodium Azide pH 7.4, then incubating the complex with 5 x 10⁵ cells at 4°C for 30 minutes. The cells were then washed three times and mean channel fluorescence at 475 nM was measured.

7. IgE In Vitro Assay Protocol

[0137]

a. Peripheral blood mononuclear cells were separated from normal donors.

b. Cells were washed extensively with phosphate buffered saline to remove as many platelets as possible.

c. Mononuclear cells were counted and resuspend in media at 1x10⁶ cells/ml. (Media=DMEM + pen/strep + 15% horse serum + IL-2 (25U/ml) + IL-4 (20ng/ml)).

d. Antibodies were added at appropriate concentrations on day 0, 5, and 8.

e. Cultures were incubated in 24 well Falcon tissue culture plates for 14 days.

f. On day 14 supernatants were removed and assayed for IgE concentrations by an IgE specific ELISA protocol.

8. Affinity constant (kd) of murine mAb for human IgE was determined by equilibrium binding (Scatchard analysis as follows:

[0138]

a. IgE (ND and PS allotypes were iodinated by the chloramine T method and separated from free ¹²⁵I Na with a PD10 sephadex G25 column (Pharmacia cat. no. 17-0851-01) in RIA buffer:PBS, 0.5% bovine serum albumin (Sigma cat. no. A-7888), 0.05% Tween 20 (Sigma cat. no. P-1379), 0.01% thimerosal (Sigma cat. no. T-5125), pH 7.4. Approximately 78-95% of the post column counts were precipitated with 50% trichloroacetic acid and specific activity of iodinated IgE preparations ranged from 1.6 to 13 µCi/µg assuming 70% counting efficiency.

b. A fixed concentration of ¹²⁵I IgE (approximately 5 x 10⁴ cpm) was added to varying concentrations of unlabelled IgE (1 to 200 nM) in a final volume of 0.1ml RIA buffer in 12x75mm polypropylene test tubes. Murine anti-human IgE mABs (20nM final concentration) in 0.1 ml RIA buffer were then added for a final assay volume of 0.2ml.

c. Samples were incubated 16-18 hours at 25°C with continuous agitation.

d. Bound and free ¹²⁵I IgE was separated by the addition of a 0.3 ml mixture of affinity purified goat anti-mouse IgG (Boehringer Mannheim cat. no. 605 208) coupled to CN Br activated Sepharose 4B (cat no. 17-0430-01) and carrier protein A sepharose (Repligen cat. no. IPA 300) in RIA buffer and incubated 1 to 2 hours at 25°C with continuous agitation. RIA buffer (1ml) was then added, and tubes were centrifuged 5 min. 400 xg. Samples were counted to determine total counts. Supernatants were aspirated with a finely drawn pasteur pipet, samples were recounted and bound versus free counts were calculated.

e. Scatchard analysis was performed utilizing a Fortran program (scanplot) based on the Ligand program written by P. Munson at NIH. Scatplot uses a mass action equation fitting bound as a function of total using the Rodbard type regression analysis.

EXAMPLE 2

Preparation of Variant IgE

[0139] Based on the model of IgE Fc by Padlan & Davies (Mol. Immunol. 23:1063 (1986), which is based on the crystal structure of human IgG1 Fc (Deisenhofer, Biochem. 20:2361-2370 [1981]), a series of mutants were designed which could be used to test the binding of human IgE to its receptors. These mutants are designated Emut 1-13, and are listed in Table 6 below. The Fc ϵ 3 domain is comprised of seven β -strands which form a β -sheet structure representative of all immunoglobulin domains; there are six loops which connect these seven β -strands. We refer to these loops by the 2 β -strands they connect, e.g. loop AB connects β -stands A and B. We have constructed mutants of human IgE in which we have substituted five of the Fc ϵ 3 domain loops with their counterparts from human IgG1 (Table 6, 1-5). The sixth loop contains the glycosylation site in both IgE and IgG and hence was not altered. One mutant, (Table 6, 6), was made by exchanging human Fc ϵ 3 β -strand D with its human IgG1 Fc γ 2 counterpart. Seven additional mutants, (Table 6, 7-13), consisted of the substitution of Ala residues into Fc ϵ 3 β -strands and a loop in Fc ϵ 2.

[0140] A human IgE gene was cloned from U266, a publicly available cell line. The gene was cloned into a previously described phagemid vector containing the human cytomegalovirus enhancer and promoter, a 5' intron and sv40 polyadenylation signal (Gorman *et al.*, DNA and Prot. Eng. Techn., 2:3-10 [1990]). Mutagenesis was performed by the Kunkel method (T.A. Kunkel, Proc. Natl. Acad. Sci. USA, 82:488-492 (1985) using buffers and enzymes supplied with the BioRad Muta-gene phagemid *in vitro* mutagenesis kit, together with oligonucleotides encoding the human IgG1 sequences shown in Table 6 below. Sequences of the mutant IgE DNAs were checked only at the site of mutation using ³⁵S dideoxy sequencing

TABLE 6

Mutant	Kabat Residue No. (Structure) ¹	Human IgE Fc ϵ 3 Seq.	Human IgG1 Fc γ 2 Seq.
1	377-385 (1AB)	FDLFIRKS (SEQ.ID NO. 10)	KDTLMISRT (SEQ.ID NO. 11)
2	396-401 (1BC)	APSKGT (SEQ.ID NO. 12)	SHEDPQ (SEQ.ID NO. 13)
3	407-420 (1CD)	SRASGKPVNHS (SEQ.ID NO. 14)	YVDGVQVHNAK (SEQ.ID NO. 15)
4	444-453 (1EF)	GTRDWIEGET (SEQ.ID NO. 16)	LHQDWLDGKE (SEQ.ID NO. 17)
5	465-469 (1FG)	RALM (SEQ.ID NO. 18)	APIE (SEQ.ID NO. 19)
6	423-428 (β D)	KEEKQR (SEQ.ID NO. 20)	PREQQY (SEQ. D NO. 21)
7	383-385 (1AB)	RKS	[AAA] ²
8	387, 389 (β B)	T(I)T	[A(I)A] ²
9	403, 405 (β C)	N(L)T	[A(L)A] ²
10	438-440 (β E)	T(S)T	[A(S)A] ²
11	455, 457, 459 (β F)	Q(C)R(V)T (SEQ.ID NO. 22)	[A(C)A(V)A] ² (SEQ.ID NO. 23)
12	471, 473 (β G)	S(T)T	[A(T)A] ²
13	329-331, 334-336	QKH(WL)SDR (SEQ.ID NO. 24)	[AAA(WL)AAA] ² (SEQ.ID NO. 25)

¹loop = 1 β -strand = β

²Sequences in brackets are from mutants in which alanine residues rather than IgG sequences were used to replace the IgE target sequence. Residues in parentheses were not altered in these mutants.

[0141] The mutant IgEs were transiently expressed in human embryonic kidney 293 cells (Gorman *et al.*, *supra*), purified on a mouse anti-human IgE antibody affinity column and samples run using SDS-PAGE to ascertain that the

mutant proteins were of the proper molecular weight.

EXAMPLE 3

5 Soluble FCEH binding assay

[0142] This assay is a sequential inhibition ELISA which measures binding to the FCEH only. In this assay, a monoclonal antibody against the FCEH is coated onto ELISA plates at a concentration of 1 µg/ml in 50 mM sodium carbonate pH 9.6 for two hours at room temperature, and blocked for two hours with PBS containing 0.5% bovine serum albumin (PBSA), then washed three times with ELISA wash buffer (0.05% Tween 20 in PBS). Recombinantly produced soluble FCEH is added at a concentration of 50 units/ml and incubated for one hour, then washed five times in ELISA wash buffer. Mutant IgE samples are then added to the wells and incubated for one to two hours. The excess mutant IgE is removed by aspiration, and biotinylated IgE is then added at 50 ng/ml for 15 minutes followed by five washes with ELISA wash buffer. Streptavidin conjugated to horseradish peroxidase (Sigma Chemical Company #S5512) was added at a 1:5000 dilution for 15 minutes, then washed three times with ELISA wash buffer. Color was developed with a tetramethyl benzidine peroxidase substrate system (Kirkegaard & Perry Labs # 50-76-00, Lot. no. NA 18) for seven minutes at 25° C. The reaction was stopped by the addition of 1 M HCl. The ability of the mutant IgE to bind the FCEH is assessed by the degree to which the biotinylated IgE is prevented from binding. This assay is designed to test for any FCEH binding by the mutant IgE and is not meant to determine the affinity of the mutant for the FCEH relative to native IgE.

FACS based binding assays for U266 IgE mutants

[0143] Tissue culture supernatants from 293s cells transfected with U266 IgE cDNA were harvested at either 48 or 96 hours post transfection. Tissue culture supernatants were concentrated 5-X with Amicon Centriprep 300 centrifugal concentrators (30,000 MW cutoff). Concentrated supernatants were passed through a mouse monoclonal anti-U266 IgE affinity column (Genentech MAE1 coupled to CnBr-Sepharose). U266 IgE was eluted from the column with 3.0 M potassium cyanate in 50 mM tris buffer Ph 7.8. Eluate fractions containing protein as determined by O.D.280 nm were pooled and placed in Amicon Centricon 300 concentrators. Eluate buffer was exchanged for PBS by passing multiple volumes of PBS through the concentrator. The final volume of affinity purified supernatant ranged from 0.5-1 ml. Structural integrity of recombinant IgE mutants was analyzed on 1-12% SDS PAGE gels and compared with U266 IgE standard obtained from the U266 cell line. Mutants were also analyzed for the ability to bind to a series of monoclonal and IgE antibodies to further ascertain proper folding and structural identity with native IgE. The concentration of immunoreactive IgE for each IgE mutant was determined by a human IgE capture ELISA as follows. Nunc Immunoplate Maxisorp plates (Nunc # 4-39451) were coated overnight at 4° C with a Genentech murine IgG1 anti-U266 IgE (MAE1) at 1 µg/ml in coat buffer (50 mM sodium carbonate buffer pH 9.6). Coat antibody was removed by three washes with ELISA wash buffer (0.05% Tween 20 (US Biochemical Corporation # 20605) in PBS). Non-specific sites were blocked with ELISA diluent buffer (50 mM tris buffered saline containing 0.5% BSA (Sigma Chemical Company # A-7888), 0.05% Tween 20 and 2 mM EDTA) for two hours at 25° C on an orbital shaker. Diluent buffer was removed with 3 washes of ELISA wash buffer. Serial two-fold dilutions of IgE mutants in ELISA diluent buffer were added to the plate. U266 IgE standard (lot 13068-46) was added at 1000, 500, 250, 125, 62.5, 31.3, and 15.6 ng/ml in duplicate as standards. Samples and standard were incubated two hours at 25° C followed by three washes with ELISA wash buffer. IgE was detected with HRP conjugated Sheep anti-human IgE (ICN # N060-050-1) at 1:8000 in ELISA diluent buffer for 90 min. at 25° C followed by 3 washes with ELISA wash buffer. HRP conjugate was developed with a tetramethyl benzidine peroxidase substrate system (Kirkegaard & Perry Labs. # 50-76-00, Lot. no. NA 18) for 7 minutes at 25° C. The reaction was stopped by the addition of 1 M HCl. The reaction product was analyzed with a dual wavelength spectrophotometer at 450 nm minus absorption at 570nm. The U266 IgE standards were used to generate a standard curve and IgE concentrations of the sample were extrapolated by non-parametric linear regression analysis.

[0144] FcER1 alpha (+) CHO 3D10 (FCEH expressing) and FcERII(CD23) (+) IM9 (FCEL expressing) B cell lines were used for the binding assays. The stably transfected CHO (duk -) cell clone 3D10 (JBC 265, 22079-22081, 1990) was maintained in Iscove's modified Dulbecco's media supplemented with 10% heat inactivated fetal calf serum, 80 µg/ml gentamicin sulfate and 5 X 10⁻⁷M methotrexate. The IM9 human B cell myeloma ATCC CCL 159. (Ann. N.Y. Acad. Sci. 190:221-234, 1972) was maintained in GIF base medium with 10% heat inactivated fetal bovine serum, penicillin, streptomycin (100 units/ml) and L-glutamine (2mM). As a positive control to determine the level of CD23 on the surface of IM9 cells in each experiment, an aliquot of cells was stained with Becton Dickinson murine monoclonal Leu 20 (anti-CD23) at 10 µg/ml for 30 minutes at 4° C followed by two washes in FACS buffer. The cells were then incubated with FITC conjugated F(ab')₂ affinity purified goat anti-murine IgG at 5 µg/ml. Adherent CHO3D10 cells were removed from tissue culture dishes by incubation with 10 mM EDTA in PBS for 2 minutes at 37°C. Cells were counted,

then resuspended in FACS buffer (0.1% BSA, 10 mM Na azide in PBS pH 7.4) at a concentration of 5×10^6 /ml. CHO3D10 and Im9 cells (5×10^5 /aliquot) were incubated in 100 μ l of FACS buffer containing U266 IgE standard or IgE mutants at 2 μ g/ml for 30 minutes at 4° C in 96 well microtiter plates followed by two washes with FACS buffer. As a control, cells were incubated in buffer alone or buffer containing 2 μ g/ml human IgG1 (Behring Diagnostics # 400112, lot no. 801024). Cells were then incubated in 100 μ l FACS buffer containing FITC conjugated rabbit anti-human IgE at 20 μ g/ml (Accurate Chem. Co. # AXL 475F, lot.no. 040A) for 30 minutes at 4° C followed by 3 washes with FACS buffer. 400 μ l of buffer containing propidium iodide at 2 μ g/ml was added to the cell suspension to stain dead cells. Cells were analyzed on a Becton Dickinson FACSCAN flow cytometer. Forward light scatter and 90 degree side scatter gates were set to analyze a homogeneous population of cells and dead cells which stained with propidium iodide were excluded from analysis. FITC positive cells (IgE binding) were analyzed relative to cells stained with FITC rabbit anti-H IgE alone.

[0145] The foregoing assays were used to determine the ability of the example 2 IgE analogues to bind to FCEH and FCEL. The results are set forth in Table 7.

TABLE 7

BINDING OF IGE AND IGE ANALOGUES TO FCEH AND FCEL			
Sample/Mutant	Conc. (ug/ml)	FCEH alpha % CHO 3D10 (+)	FCEL (CD23) % IM9 (+)
U266 IgE	10	90.3	92.5
U266 IgE	5	89.9	82.6
U266 IgE	0.5	59.6	4.6
U266 IgE	0.1	15.8	1.7
7	1.65 ¹	1.7	4.3
2	1.65	34.3	48.9
3	1.65	32.3	1.2
4	1.65	4.9	9.2
5	1.65	60.5	73.9
6	1.65	1.4	71.6
7	1.65	76.4	4.6
8	1.65	70.3	16.3
3	1.65	84.2	94.3
10	1.65	67.5	84.8
11	1.65	70.8	61.5
12	1.65	84.7	90.3
13	1.65	85.7	96.1
dh 184 (+)	1.65	83.8	21.1
PA13 ² (control)	10	1.3	

¹Values based on quantitative Elisa. U266 was used as the standard and murine anti-F α monoclonal antibody to capture.

²A CDR grafted human IgG.

[0146] Three mutant IgEs exhibited complete loss of binding to the FCEH receptor: mutants 1, 4 and 6. Mutant 6 altered β -strand D at the end of Fc ϵ 3 close to the Fc ϵ 2 domain. Mutants 1 and 4 involved alteration of two Fc ϵ 3 loops which are adjacent and near the Fc ϵ 4 domain. Note that mutant 7 is subset of mutant 1 in which the three C-terminal residues of loop AB have been changed to alanines (Table 6, 1 vs. 7). However, mutant 7 does not affect binding to FCEH. We interpret this to mean that either 1) Fc ϵ RI binds at least one of IgE residues 377-381 or 2) the extra residue in IgG1 loop AB (9 residues) substituted for IgE loop AB (8 residues) effected deformation of some adjacent binding determinant, possibly loop EF. That mutants 8 and 10 had no affect on Fc ϵ RI binding most likely means that the FCEH receptor does not protrude into the cavity bounded by loop AB and β -strand D.

[0147] Although mutant 4 had a Leu replacing Gly444 (Table 6), this should not affect the conformation of loop EF.

Residue 444 is prior to the N-terminus of this α -helix. In addition, murine IgE has a Val at position 444 and rat IgE has an Asp. The two buried hydrophobic residues in the middle of the α -helix, W448 and I449, are retained in the substituted IgG1 loop (W448, L449) as is G451 which terminated the α -helix. Hence the conformation of loop EF should be similar in IgE and IgG1.

[0148] Mutants 2 and 3 exhibited decreased binding to FCEH. Since loop BC lies near β -strand D and loop CD is in the vicinity of loop EF, it is conceivable that one or two residues in loops BC and CD contact FCEH.

[0149] Five mutant IgEs exhibited loss of binding to the FCEL receptor: mutants 1, 3, 4, 7 and 8. Mutants 1 and 4 were discussed above. Mutant 3 involved alteration of loop CD; in contrast to FCEH, loop CD evidently plays a major role in FCEL binding. Mutant 7, a subset of mutant 1 as discussed above, comprises the C-terminal portion of loop AB and is proximal to loop EF. Additionally, mutant 8 consists of replacement of two Thr residues (387,389) with Ala; these two residues are part of β -strand B which is at the bottom of the aforementioned cavity bounded by loop AB and β -strand D. Mutant 10 comprised a different two residues in this cavity (438,440) on β -strand E, which is adjacent to β -strand B. Since mutant 10 did not affect FCEL binding, we conclude that the FCEL receptor should have only a minimal incursion into cavity while the high affinity receptor does not intrude into the cavity.

[0150] In addition to a glycosylation site at Asn430 which corresponds to the glycosylation site in IgG Fc, human IgE contains another glycosylation site at Asn403. Mutant 9 converted Asn403 and Thr405 to alanines (Table 6). Loss of carbohydrate did not affect binding to either receptor.

[0151] Based on the information from mutants 1-13, we propose that FCEH and FCEL have binding sites on IgE Fc which are distinct but overlap. The low affinity receptor seems to interact with a relatively smaller portion of the IgE Fc ϵ 3 domain involving three adjacent loops: AB, CD and EF. In contrast, the high affinity receptor interacts with a larger portion of IgE Fc ϵ 3, which spans loop EF, β -strand D and, possibly, the N-terminal portion of loop AB. Portions of loops BC and CD in the vicinity of loop EF and β -strand D may also interact with FCEH. In addition, FCEL may protrude into the cavity bounded by loop AB and β -strand D whereas FCEH does not do so. Since we have not evaluated any mutants in Fc ϵ 4 and only one in Fc ϵ 2 (mutant 13), it is possible that portions of these two domains play a role in IgE-receptor binding.

EXAMPLE 4

Preparation of Humanized MaE11

[0152] Residues were selected from MaE11 and inserted or substituted into a human Fab antibody background (V_H region Kabat subgroup III and V_L region kappa subgroup I). A first version, humaellvl or version 1, is described in Table 8.

TABLE 8.

Changes in V_H human subgroup III and V_L κ subgroup I (Kabat) consensus sequences for humanized MaE11 Version 1

Domain	hu Residue	Residue No.	V.1	CDR by Rabat	CDR by Chothia
V _L					
	M	4	L		
	insert	30abcd	YDGD (SEQ.ID. NO. 26)	L1	L1
	L*	33	M	L1	
	S	53	Y	L2	
	Y	31	S	L3	L3
	N	92	H	L3	L3
	S	93	E	L3	L3
	L	94	D	L3	L3
V _H					
	A	24	V		
	L*	27	Y	H1	H1
	T	28	S	H1	H1
	F*	29	I	H1	H1
	insert	29a	T	H1	H1
	D	31	G	H1	H1
	A	33	S	H1	H1
	M*	31	W	H1	H1
	V	37	I		
	V	50	S	H2	
	S	58	I	H2	
	N	53	Y	H2	H2
	D	54	D	H2	H2
	S	50	G	H2	H2
	Y	58	N	H2	
	L	78	F		
	D	95	G	H3	
		97-101	All Changed to MaE11 Sequence	H3	H3

* These residues typically do not vary despite their position within CDRs. The remaining residues found in the KI and III CDR sequences (particularly the CDRs by Chothia structural analysis), will vary widely among recipient human antibodies.

[0153] The affinity of version 1 was assayed and found to be about 100 times lower than that of the donor antibody Mae11 (see Figs. 4a and 4b). Therefore, further modifications in the sequence of version 1 were made as shown in Table 9. Determination was made of the ability of these further modifications to inhibit the binding of labelled hulGE to FCEH.

[0154] The 50% inhibition assays whose results are shown in Table 9 were conducted as follows:

[0155] A 96-well assay plate (Manufn Nunc.) was coated with 0.05 ml of the Fc ϵ RI alpha chain IgG1 chimeric receptor in 1 μ g/ml coating buffer (SONmol carbonate/bicarbonate, pH 9.6). Assay was done for 12 hours at 4-8° C. The wells were aspirated and 250 μ l blocking buffer (PBS--1% BSA pH 7.2) was added and incubated for one hour at 4°C. In a separate assay plate the samples and reference murine MaE11 antibody were titrated from 200 μ g/ml by 1 to 10-fold dilution with assay buffer (0.5% BSA, 0.05% Tween 20, PBS, pH 7.2) and an equal volume of 10ng/ml biotinylated IgE at 10ng/ml was added and the plate incubated for 2-3 hours at 25°C. The Fc ϵ RI-coated wells were washed three times with PBS-0.05% Tween20, and then 50 μ l from the sample wells were transferred and incubated with agitation for 30 minutes at 25°C. 50 μ l/well of streptavidin-HRP diluted 1:5000 in assay buffer was incubated for 15 minutes with agitation and then the plate was washed as before. 50 μ l/well of Microwell peroxidase substrate (Kirkgaard & Parry Laboratories) was added and color was developed for 30 minutes. The reaction was stopped by adding an equal volume of 1 normal HCl and the adsorbance measured at 450nm. The concentration for 50% inhibition was calculated by plotting percent inhibition versus concentration of blocking antibody with a nonlinear 4-parameter curve-fit for each antibody using INPLOT.

TABLE 9

Humanized MaE11 Variants						
Version [F(ab)-X]	Domain	Changes from F(ab)-Version 1	Purpose	Conc. at 60% Inh. (ng/ml)* Mean	S.D. for prev. col.	F(ab)-X / F(ab)-1
1	-	-	-	6083	1279	1.0
2	V _L	L4M M33L	Packing; CDR-L1	9439	508	1.6
3	V _L	E55G G57E	Sequence usually E55-X-G57	5799	523	1.0
4	V _H	I37V	VL-VH interface	8622	107	1.4
5	V _H	V24A	Packing; CDR-H1	9387	733	1.6
6	V _H	F78L	Packing; CDR-H1,H2	17537	4372	2.9
7	V _L V _H	L4M R24K E55G G57E V24A I37V T57S A60N D61P V63L G65N F78L	remake version 1 to accomplish a direct exchange of CDR residues	> 100000		> 16.0#
7a	V _H	As V.7 except V _H L78 is F		98000		16.0

* Inhibition of flic-IgE binding to FcεRH (FcεRI). Full length antibody and humanized fragment versions: mean and standard deviation of three assays.
 # A F(ab)-X / F(ab)-1 ratio of > 16 means that this variant exhibited no binding even at the highest F(ab) concentrations used.

TABLE 9 (continued)

Humanized MaE11 Variants						F(ab)-X
Version [F(ab)-X]	Domain	Changes from F(ab)-Version 1	Purpose	Conc. at 60% Inh. (ng/ml)* Mean	S.D. for prev. col.	F(ab)-1
8	V _H	A60N D61P	Extended Kabat CDR-H2 & A60N is at V _L -V _H interface	1224	102	0.20
8a	V _H	As V.8 except V _H V62 is L and F67 is I	CDR-H2; packing of L63 and I67	416	66	0.07
8b	V _H	As V.8 except F67 is I	CDR-H2; packing of V63 and I67	501	84	0.08
1	-	-	-	6083	1279	1.0
9	V _L	A13V V19A V58I L78V V104L V48M A49G A60N V63L F67I I69V M82L L82cA	Repack Version 1 interior as in murine MaE11	842	130	0.14
23	V _L	L4M	Packing; CDR-L1	6770	349	1.1
10	V _L	D29aA D29cA D30A	CDR-L1 modification	>100000		>16.0
11	V _L	E93A D94A	CDR-L3 modification	17456	7115	2.9

* Inhibition of flic-IgE binding to FcEH (FcERI). Full length antibody and humanized fragment versions: mean and standard deviation of three assays.

TABLE 9 (continued)

Humanized MaE11 Variants						F(ab)-X
Version [F(ab)-X]	Domain	Changes from F(ab)-Version 1	Purpose	Conc. at 60% inh. (ng/ml)* Mean	S.D. for prev. col.	F(ab)-1
12	V _H	D54A	CDR-H2 modification	2066	174	0.34
13	V _H	H97A H100aA H100cA	CDR-H3 modification	>100000		> 16.0
14	V _L	D29aA	CDR-L1 modification	3452	183	0.57
15	V _L	D29cA	CDR-L1 modification	6384	367	1.0
16	V _L	D30A	CDR-L1 modification	>100000		> 16.0
17	V _H	H97A	CDR-H3 modification	19427	8360	3.2
18	V _H	H100aA	CDR-H3 modification	2713	174	0.45
19	V _H	H100cA	CDR-H3 modification	15846	8128	2.6

* Inhibition of fltc-IgE binding to FcEH (FcERI). Full length antibody and humanized fragment versions: mean and standard deviation of three assays.

[0156] As can be seen from Table 9 and Figs. 4a and 4b, version 8 (in which human residues of version 1 at sites 60 and 61 in the light chain were replaced by their Mae11 counterparts) demonstrated substantially increased affinity. Further increases in affinity are seen in versions 8a and 8b, where one or two murine residues replaced human residues. Other increases, at least virtually to the level of Mae11, were accomplished by replacing hydrophobic human residues found in the interior of V_H and V_{H1} with their Mae11 counterparts, resulting in the variant designated version 9 (see Table 9 and Figs. 4a and 4b). Accordingly, the humanized antibodies of this invention will possess affinities ranging about from 0.1 to 100 times that of MAE11.

[0157] Table 10 explores the effects on FCEH affinity of various combinations of humanized maE11 IgG1 variants.

Table 10.

Humanized MaE11 IgG1 Variants				
Variant	Conc. at 50% inh. (ng/ml) Mean*	S.D. from previous column*	Var. X IgL1H1	Var. X MaE11
IgL1H1	7569	1042	1.0	16.9
IgL1H8	3493	1264	0.46	7.8
IgL9H9	1118	172	0.15	2.5
IgL1H9	608	364	0.08	1.4
IgL9H1	5273	2326	0.70	11.7
IgL1H8b	1449	226	0.19	3.2
MaE11	449	53	0.06	1.0

* L1 = V_L as in F(ab)-1 (human buried residues--not exposed to solvent); L9 = V_L as in F(ab)-9 (murine buried residues); H1 = V_H as in F(ab)-1 (human buried residues); H8 = V_H as in F(ab)-8 (F(ab)-1 with AlaH60Asn, AspH61Pro); H9 = V_H as in F(ab)-9 (murine buried residues); H8b = V_H as in F(ab)-8b (F(ab)-8 with PheH67Ile).

EXAMPLE 5

Creation of IgE Mutants

[0158] IgE mutants (Table 11) were prepared to evaluate their effect on binding to anti-IgE, especially MaE11, and to Fc ϵ RI and Fc ϵ RII. Some of the mutants were designed to substitute for a specific amino acid residue another residue with either similar or very different charge or size. The impact of these changes on receptor binding is reflected in the table below.

[0159] The receptor assays are performed substantially as follows:

[0160] A 96-well assay plate (Manufn Nunc.) was coated with 0.05 ml of Fc ϵ RI or RII IgG1 chimeric receptor in 1 μ g/ml coating buffer (50nmol carbonate/bicarbonate, pH 9.6). Assay was done for 12 hours at 4-8° C. The wells were aspirated and 250 μ l blocking buffer (PBS--1% BSA pH 7.2) was added and incubated for one hour at 4°C. In a separate assay plate the samples and reference murine MaE11 antibody were titrated from 200 μ g/ml by 1 to 10-fold dilution with assay buffer (0.5% BSA, 0.05% Tween 20, PBS, pH 7.2) and an equal volume of 10ng/ml biotinylated IgE at 10ng/ml was added and the plate incubated for 2-3 hours at 25°C. The Fc ϵ RI-coated wells were washed three times with PBS-0.05% Tween20, and then 50 μ l from the sample wells were transferred and incubated with agitation for 30 minutes at 25°C. 50 μ l/well of streptavidin-HRP diluted 1:5000 in assay buffer was incubated for 15 minutes with agitation and then the plate was washed as before. 50 μ l/well of Microwell peroxidase substrate (Kirkgaard & Parry Laboratories) was added and color was developed for 30 minutes. The reaction was stopped by adding an equal volume of 1 normal HCl and the adsorbance measured at 450nm. The absorbance was plotted versus concentration of blocking antibody MaE11 and an inhibition standard curve was generated using INPLOT.

Table 11.

Amino acid sequences of IgE mutants					
Mutant	Kabat residue #	Human IgE Fce3 seq.	Mutant seq.	Fcε-RI*	FcεRII *
Loop AB					
1	377-385	FDLFIRKS (SEQ.ID.27)	KDTLMISRT (SEQ.ID.28)	-	-
7	383-385	RKS	AAA	+/-,-	+,-
21	377, 381	F(DL)F (SEQ.ID.29)	Q(DL)H (SEQ.ID.30)	+	+
66	382	I	A	+	+
67	383	R	A	+	+/-
68	384	K	A	+	+
69	385	S	A		
102	383, 384	RK	DD		
β-strand B					
8	387, 389	T(I)T	A(I)A	+/-,+	-
70	387	T	A	+	+/-,+
71	389	T	A	+	+
Loop BC					
2	396-401	APSKGT (SEQ.ID.31)	SHEDPQ (SEQ.ID.32)		
β-strand C					
9	403, 405	N(L)T	A(L)A	+	+
Loop CD					
3	407-420	SRASGKPVNHS (SEQ.ID.33)	YVDGVQVHNAK (SEQ.ID.34)	+/-	-
55	407-415	SR(A)S(G)K (SEQ.ID.35)	AA(A)A(G)A (SEQ.ID.36)	+/-	+
59	407	S	A	+	+
60	408	R	A	+	-
61	411	S	A	+	+
62	415	K	A	+	-
63	418	N	A	+/-	+
64	419	H	A	+	+
65	420	S	A	+/-	+
100	408	R	E		
101	415	K	D		
β-strand D					
6	423-428	KEEKQR (SEQ.ID.37)	PREQQY (SEQ.ID.38)	+	+
35	422	R	A	+	+
36	4423	K	A	+	+
37	424	E	A	+	+
38	425	E	A	+	+
39	426	K	A	+	

* Positive receptor binding indicated by "+", no binding by "-", and positive binding but less than unaltered is shown by "+/-". Where more than one assay was performed, results are separated by commas.

Table 11. (continued)

Amino acid sequences of IgE mutants					
Mutant	Kabat residue #	Human IgE Fcε3 seq.	Mutant seq.	Fcε-RI*	FcεRII *
β-strand D					
40	427	Q	A	-,+/-	+
41	428	R	A	+	
75	423-425	KEE	AAA	-,+/-,+	+
76	426-428	KQR	AAA		
79	423,425,427	KEEKQR (SEQ.ID.39)	AEAKAR (SEQ.ID.40)		
80	424,426,428	KEEKQR (SEQ.ID.41)	KAEAQA (SEQ.ID.42)		
81	28	K	P		
82	423,423-427	KEEKQR (SEQ.ID.43)	AAEAQA (SEQ.ID.44)		
β-strand E					
10	438,440	T(S)T	A(S)A	+	+
Loop EF					
4	444-453	GTRDWIEGET (SEQ.ID.45)	LHQDWLDGKE (SEQ.ID.46)	-	-
49	445	T	A	+	+
50	336	R	A	+	-
51	337	D	A	+	+,-
52	450	E	A	+/-	
53	452	E	A	+	-
77	445,446	TR	AA	+	+
78	450,452,453	E(G)ET (SEQ.ID.47)	A(G)AA (SEQ.ID.48)	+	+
83		G	L	+	+
84	444	G	A	+	+
85	444	TRDWIEGET (SEQ.ID.49)	HQDWLDGKE (SEQ.ID.50)	-	+
86	445	T	H	+	
87	445	TR	HQ	+	
88	445,446	R	E	+/-,+	
89	446	E(G)ET (SEQ.ID.51)	D(G)KE (SEQ.ID.52)	-	
93	450,452,453	D	R	+/-,-	+/-
94		E	R	+	
95	447	E	R		
96	450	T	R		
97	452	D	N		
98	453	E	Q		
99	447	E	D		
	452				

* Positive receptor binding indicated by "+", no binding by "-", and positive binding but less than unaltered is shown by "+/-". Where more than one assay was performed, results are separated by commas.

Table 11. (continued)

Amino acid sequences of IgE mutants					
Mutant	Kabat residue #	Human IgE Fcε3 seq.	Mutant seq.	Fcε-R1*	Fcε-R2 *
Loop EF					
	452				
β-strand F					
11	445,457,459	Q(C)R(V)T (SEQ.ID.53)	A(C)A(V)A (SEQ.ID.54)		
Loop FG					
5	465-469	RALM (SEQ.ID.55)	APIE (SEQ.ID.56)		
β-strand G					
12	471,473	S(T)T	A(T)A	+,+	
Fcε2					
13	329-331, 334-336	QKH(WL)SDR (SEQ.ID.57)	AAA(WL)AAA (SEQ.ID.58)	+,+	
Fcε4					
72	498-501	PRAA (SEQ.ID.59)	QPRE (SEQ.ID.60)		
73	594-599	ASPSQT (SEQ.ID.61)	LHNHY (SEQ.ID.62)		
74	595-599	S(P)SQT (SEQ.ID.63)	A(P)AAA (SEQ.ID.64)		

* Positive receptor binding indicated by "+", no binding by "-", and positive binding but less than unaltered is shown by "+/-". Where more than one assay was performed, results are separated by commas.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Jardieu, Paula M.
Presta, Leonard G.

(ii) TITLE OF INVENTION: Immunoglobulin Variants

(iii) NUMBER OF SEQUENCES: 64

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Genentech, Inc.
(B) STREET: 460 Point San Bruno Blvd
(C) CITY: South San Francisco
(D) STATE: California
(E) COUNTRY: USA
(F) ZIP: 94080

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: 5.25 inch, 360 Kb floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: patin (Genentech)

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE: 14-AUG-1992
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 07/879495
(B) APPLICATION DATE: 07-MAY-1992

(viii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 07/744768
(B) APPLICATION DATE: 14-AUG-1991

(ix) ATTORNEY/AGENT INFORMATION:

(A) NAME: Adler, Carolyn R.
(B) REGISTRATION NUMBER: 32,324
(C) REFERENCE/DOCKET NUMBER: 718P2

(x) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 415/225-2614
(B) TELEFAX: 415/952-9881
(C) TELEX: 910/371-7168

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 118 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Xaa	Asp	Ser	Asn	Pro	Arg	Gly	Val	Ser	Ala	Tyr	Leu	Ser	Arg	Pro
1				5					10					15
Ser	Pro	Phe	Asp	Xaa	Leu	Phe	Ile	Arg	Lys	Ser	Pro	Thr	Ile	Thr
				20					25					30
Cys	Leu	Val	Val	Asp	Leu	Ala	Pro	Ser	Lys	Gly	Thr	Val	Asn	Leu
				35					40					45
Thr	Trp	Ser	Arg	Xaa	Ala	Ser	Xaa	Xaa	Gly	Lys	Pro	Val	Asn	His

50 55 60

Ser Thr Arg Lys Glu Glu Lys Gln Arg Xaa Asn Xaa Xaa Gly Thr
 65 70 75

Leu Thr Val Thr Ser Thr Leu Pro Val Gly Thr Arg Asp Trp Ile
 80 85 90

Glu Gly Glu Thr Gln Cys Arg Val Thr His Pro His Leu Pro Arg
 95 100 105

Ala Leu Xaa Met Arg Ser Thr Thr Lys Thr Ser Gly Pro
 110 115 118

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 111 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu
 1 5 10 15

Gly Gln Arg Ala Thr Ile Ser Cys Lys Ala Ser Gln Ser Val Asp
 20 25 30

Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly
 35 40 45

Gln Pro Pro Ile Leu Leu Ile Tyr Ala Ala Ser Tyr Leu Gly Ser
 50 55 60

Glu Ile Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe
 65 70 75

Thr Leu Asn Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Phe
 80 85 90

Tyr Cys Gln Gln Ser His Glu Asp Pro Tyr Thr Phe Gly Ala Gly
 95 100 105

Thr Lys Leu Glu Ile Lys
 110 111

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 134 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Asp Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser
 1 5 10 15

Gln Ser Leu Ser Leu Ala Cys Ser Val Thr Gly Tyr Ser Ile Thr
 20 25 30

Ser Gly Tyr Ser Trp Asn Trp Ile Arg Gln Phe Pro Gly Asn Lys
 35 40 45

Leu Glu Trp Met Gly Ser Ile Thr Tyr Asp Gly Ser Ser Asn Tyr
 50 55 60

Asn Pro Ser Leu Lys Asn Arg Ile Ser Val Thr Arg Asp Thr Ser
 65 70 75
 5 Gln Asn Gln Phe Phe Leu Lys Leu Asn Ser Ala Thr Ala Glu Asp
 80 85 90
 Thr Ala Thr Tyr Tyr Cys Ala Arg Gly Ser His Tyr Phe Gly His
 95 100 105
 10 Trp His Phe Ala Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser
 110 115 120
 Ser Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala Arg
 125 130 134

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 124 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Asp Ile Val Met Thr Gln Ser Gln Lys Phe Met Ser Thr Ser Val
 1 5 10 15
 Gly Asp Arg Val Ser Val Thr Cys Lys Ala Ser Gln Asn Val Ser
 25 20 25 30
 Ser Asn Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys
 35 40 45
 Ala Leu Ile Tyr Ser Ala Ser Tyr Arg Tyr Ser Gly Val Pro Asp
 30 50 55 60
 Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
 65 70 75
 Ser Asn Val Gln Ser Glu Asp Leu Ala Glu Tyr Phe Cys Gln Gln
 80 85 90
 35 Tyr Tyr Thr Tyr Pro Leu Tyr Thr Phe Gly Gly Gly Thr Lys Leu
 95 100 105
 Glu Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro
 110 115 120
 40 Pro Ser Thr Arg
 124

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 130 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Asp Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser
 1 5 10 15
 Gln Ser Leu Ser Leu Thr Cys Thr Val Thr Gly Tyr Thr Ile Thr
 20 25 30
 55 Ser Asp Asn Ala Trp Asn Trp Ile Arg Gln Phe Pro Gly Asn Lys

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 106 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

[illegible]

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 137 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

Asp	Val	Gln	His	Gln	Glu	Ser	Glu	Pro	Asp	Leu	Val	Lys	Pro	Ser
1				5					10					15
Gln	Ser	Leu	Ser	Leu	Thr	Cys	Thr	Val	Thr	Gly	Tyr	Ser	Ile	Thr
				20					25					30

Ser Gly Tyr Asn Arg His Trp Ile Arg Gln Phe Pro Gly Asn Lys
 35 40 45
 5 Leu Glu Trp Met Gly Tyr Ile His Tyr Ser Gly Ser Thr Asn Tyr
 50 55 60
 Asn Pro Ser Leu Lys Arg Arg Ile Ser Ile Thr Arg Asp Thr Ser
 65 70 75
 10 Lys Asn Gln Phe Phe Leu Gln Leu Asn Ser Val Thr Thr Glu Asp
 80 85 90
 Thr Ala Thr Tyr Tyr Cys Ala Arg Gly Ser Ile Tyr Tyr Tyr Gly
 95 100 105
 15 Ser Arg Tyr Arg Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr Val
 110 115 120
 Thr Val Ser Ser Ala Lys Arg His Pro His Leu Ser Ile His Trp
 125 130 135
 20 Pro Gly
 137

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 453 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15
 30 Gly Ser Leu Arg Leu Ser Cys Ala Val Ser Gly Tyr Ser Ile Thr
 20 25 30
 Ser Gly Tyr Ser Trp Asn Trp Ile Arg Gln Ala Pro Gly Lys Gly
 35 35 40 45
 Leu Glu Trp Val Ala Ser Ile Thr Tyr Asp Gly Ser Thr Asn Tyr
 50 55 60
 Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser
 65 70 75
 40 Lys Asn Thr Phe Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 80 85 90
 Thr Ala Val Tyr Tyr Cys Ala Arg Gly Ser His Tyr Phe Gly His
 95 100 105
 45 Trp His Phe Ala Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 110 115 120
 Ser Ala Ser Thr Lys Gly Lys Gly Pro Ser Val Phe Pro Leu Ala
 125 130 135
 50 Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
 140 145 150
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 155 160 165
 55 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu

	170	175	180
5	Gln Ser Ser Gly Leu Tyr Ser Leu Ser	Ser Val Val Thr Val Pro	
	185	190	195
	Ser Ser Ser Leu Gly Thr Gln Thr Tyr	Ile Cys Asn Val Asn His	
	200	205	210
10	Lys Pro Ser Asn Thr Lys Val Asp Lys	Lys Val Glu Pro Lys Ser	
	215	220	225
	Cys Asp Lys Thr His Thr Cys Pro Pro	Cys Pro Ala Pro Glu Leu	
	230	235	240
	Leu Gly Gly Pro Ser Val Phe Leu Phe	Pro Pro Lys Pro Lys Asp	
	245	250	255
15	Thr Leu Met Ile Ser Arg Thr Pro Glu	Val Thr Cys Val Val Val	
	260	265	270
	Asp Val Ser His Glu Asp Pro Glu Val	Lys Phe Asn Trp Tyr Val	
	275	280	285
20	Asp Gly Val Glu Val His Asn Ala Lys	Thr Lys Pro Arg Glu Glu	
	290	295	300
	Gln Tyr Asn Ser Thr Tyr Arg Val Val	Ser Val Leu Thr Val Leu	
	305	310	315
25	His Gln Asp Trp Leu Asn Gly Lys Glu	Tyr Lys Cys Lys Val Ser	
	320	325	330
	Asn Lys Ala Leu Pro Ala Pro Ile Glu	Lys Thr Ile Ser Lys Ala	
	335	340	345
30	Lys Gly Gln Pro Arg Glu Pro Gln Val	Tyr Thr Leu Pro Pro Ser	
	350	355	360
	Arg Glu Glu Met Thr Lys Asn Gln Val	Ser Leu Thr Cys Leu Val	
	365	370	375
35	Lys Gly Phe Tyr Pro Ser Asp Ile Ala	Val Glu Trp Glu Ser Asn	
	380	385	390
	Gly Gln Pro Glu Asn Asn Tyr Lys Thr	Thr Pro Pro Val Leu Asp	
	395	400	405
40	Ser Asp Gly Ser Phe Phe Leu Tyr Ser	Lys Leu Thr Val Asp Lys	
	410	415	420
	Ser Arg Trp Gln Gln Gly Asn Val Phe	Ser Cys Ser Val Met His	
	425	430	435
45	Glu Ala Leu His Asn His Tyr Thr Gln	Lys Ser Leu Ser Leu Ser	
	440	445	450
	Pro Gly Lys		
	453		

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 218 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Val Asp
 5 20 25 30
 Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly
 35 40 45
 Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Tyr Leu Glu Ser
 10 50 55 60
 Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe
 65 70 75
 Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr
 15 80 85 90
 Tyr Cys Gln Gln Ser His Glu Asp Pro Tyr Thr Phe Gly Gln Gly
 95 100 105
 Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe
 20 110 115 120
 Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser
 125 130 135
 Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val
 25 140 145 150
 Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
 155 160 165
 Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 170 175 180
 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
 30 185 190 195
 Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr
 200 205 210
 Lys Ser Phe Asn Arg Gly Glu Cys
 35 215 218

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 8 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Phe Asp Leu Phe Ile Arg Lys Ser
 1 5 8

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Lys Asp Thr Leu Met Ile Ser Arg Thr

1 **5** **9**

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SBQ ID NO:12:

Ala Pro Ser Lys Gly Thr
1 5 6

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Ser His Glu Asp Pro Gln
1 5 6

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ser Arg Ala Ser Gly Lys Pro Val Asn His Ser
1 5 10 11

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Tyr Val Asp Gly Val Gln Val His Asn Gln Lys
1 5 10 11

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Gly Thr Arg Asp Trp Ile Glu Gly Glu Thr
1 5 10

(2) INFORMATION FOR SEQ ID NO:17:

(1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids
(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Leu His Gln Asp Trp Leu Asp Gly Lys Glu
1 5 10

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Arg Ala Leu Met
1 4

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Ala Pro Ile Glu
1 4

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Lys Glu Glu Lys Gln Arg
1 5 6

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Pro Arg Glu Gln Gln Tyr
1 5 6

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Gln Cys Arg Val Thr
1 5

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Ala Cys Ala Val Ala
 1 5

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 8 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Gln Lys His Trp Leu Ser Asp Arg
 1 5 8

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 8 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Ala Ala Ala Trp Leu Ala Ala Ala
 1 5 8

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Tyr Asp Gly Asp
 1 4

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 8 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Phe Asp Leu Phe Ile Arg Lys Ser
 1 5 8

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Lys Asp Thr Leu Met Ile Ser Arg Thr
1 5 9

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Phe Asp Leu Phe
1 4

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Gln Asp Leu His
1 4

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Ala Pro Ser Lys Gly Thr
1 5 6

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Ser His Glu Asp Pro Gln
1 5 6

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Ser Arg Ala Ser Gly Lys Pro Val Asn His Ser
1 5 10 11

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Tyr Val Asp Gly Val Gln Val His Asn Ala Lys
 1 5 10 11

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Ser Arg Ala Ser Gly Lys
 1 5 6

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Ala Ala Ala Ala Gly Ala
 1 5 6

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Lys Glu Glu Lys Gln Arg
 1 5 6

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

Pro Arg Glu Gln Gln Tyr
 1 5 6

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

Lys Glu Glu Lys Gln Arg
1 5 6

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

Ala Glu Ala Lys Ala Arg
1 5 6

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Lys Glu Glu Lys Gln Arg
1 5 6

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

Lys Ala Glu Ala Gln Ala
1 5 6

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

Lys Glu Glu Lys Gln Arg
1 5 6

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

Ala Ala Glu Ala Gln Ala
1 5 6

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids

(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

Gly Thr Arg Asp Trp Ile Glu Gly Glu Thr
1 5 10

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Leu His Gln Asp Trp Leu Asp Gly Lys Glu
1 5 10

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Glu Gly Glu Thr
1 4

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

Ala Gly Ala Ala
1 4

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

Thr Arg Asp Trp Ile Glu Gly Glu Thr
1 5 9

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

His Gln Asp Trp Leu Asp Gly Lys Glu
1 5 9

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

Glu Gly Glu Thr
 1 4

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

Asp Gly Lys Glu
 1 4

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

Gln Cys Arg Val Thr
 1 5

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

Ala Cys Ala Val Ala
 1 5

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Arg Ala Leu Met
 1 4

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

Ala Pro Ile Glu
1 4

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Gln Lys His Trp Leu Ser Asp Arg
1 5 8

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Ala Ala Ala Trp Leu Ala Ala Ala
1 5 8

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

Pro Arg Ala Ala
1 4

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

Gln Pro Arg Glu
1 4

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Ala Ser Pro Ser Gln Thr
1 5 6

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

Leu His Asn His Tyr
 1 5

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

Ser Pro Ser Gln Thr
 1 5

(2) INFORMATION FOR SEQ ID NO:64:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

Ala Pro Ala Ala Ala
 1 5

Claims

1. An antibody which is capable of binding to Fc ϵ RII bound IgE but is substantially incapable of binding to Fc ϵ RI bound IgE.
2. The antibody according to claim 1 comprising a human Kabat CDR domain into which has been substituted an analogous residue from a Kabat CDR domain of MAE11, MAE13, MAE15 or MAE17.
3. The antibody according to claim 2, wherein the residue is from the MAE11, MAE13 or MAE15 Kabat VH1 CDR domain.
4. The antibody according to claim 2 or claim 3, wherein the substituted amino acid sequence comprises from 1 to 7 residues from a MAE11, MAE13 or MAE15 Kabat CDR domain.
5. The antibody according to any one of the claims 2 to 4, wherein the substituted residue is from MAE11, MAE13 or MAE15 Kabat VH1, VH2, VH3, VL1, VL2 and VL3 domains.
6. The antibody according to any one of the claims 2 to 5, which comprises non-CDR sequence from a Kabat human consensus antibody.
7. The antibody according to claim 6, wherein the consensus antibody is Kabat subgroup III for heavy chain and kappa subgroup I for light chain.
8. The antibody according to any one of claims 2 to 7, further comprising a residue substituted from a MAE11, MAE13, MAE15 or MAE17 framework or VH-VL interface domain into the analogous residue of the human antibody.
9. The antibody according to any one of claims 2 to 8, wherein the residue is from the heavy chain framework.

10. The antibody according to claim 9, wherein the residue is VH78, VH60 or VH61.
11. An antibody which is capable of binding to Fc ϵ RII-bound IgE but is substantially incapable of binding to Fc ϵ RI-bound IgE, comprising the heavy and light chain sequences of humaellver. 1, 2, 3, 4, 5, 6, 7, 7a, 8, 8a, 8b or 9.
12. The antibody according to claim 11 which is humaellver.9.
13. The antibody according to any one of the preceding claims, wherein the antibody is bispecific.
14. An antibody which is (a) monovalent for Fc ϵ RII-bound IgE but is substantially incapable of binding to Fc ϵ RI-bound IgE and (b) is capable of an immunoglobulin effector function and comprises an FC domain containing at least two heavy chains.
15. An antibody which is capable of binding to FcRII-bound IgE but is substantially incapable of binding to FcRI-bound IgE, comprising a human consensus heavy chain and light chain sequence.
16. The antibody according to claim 15, wherein the consensus heavy chain is Kabat subgroup III and the consensus light chain is Kabat kappa subgroup I.
17. An antibody which is capable of binding to FcRII-bound IgE is substantially incapable of binding to FcRI-bound IgE, comprising a human heavy chain and light chain sequence, and which has an IgE affinity which is substantially the same as or greater than that of MAE11 for IgE.
18. The antibody according to claim 17, wherein the affinity for IgE is about 0.1 to 100 times greater than that of MAE11 for IgE.
19. The antibody according to claim 17 or claim 18, wherein the human heavy chain or light chain sequence comprises a residue substituted from MAE11, MAE13 or MAE15.
20. The antibody according to claim 1 wherein the antibody is an immunoglobulin.
21. The immunoglobulin according to claim 20 which is an IgE analogue having a variant amino acid sequence within about residues 420 to 428, inclusive.
22. The immunoglobulin according to claim 20 or claim 21 which is an IgE analogue having a variant amino acid sequence within about residues 446 to 453, inclusive.
23. The immunoglobulin according to claim 21, further comprising IgE residues about from 373-390 and wherein the variant amino acid sequence is a deletion of one of residues 423-428.
24. The immunoglobulin according to any one of claims 20 to 23 which further comprises a cytotoxic polypeptide, an enzyme, a diagnostic label, or an immunoglobulin variable domain capable of binding predetermined antigen.
25. The immunoglobulin according to claim 20 which is an IgE analogue having a variant amino acid sequence within about residues 420-428, inclusive, and within about residues 446-453, inclusive.
26. The immunoglobulin according to claim 25 which is capable of binding complement.
27. The immunoglobulin according to any one of claim 24 to 26, wherein the antigen is CD8 or CD3.
28. The immunoglobulin according to any one of claims 24 to 26, wherein the antigen is a lymphoid cell surface antigen.
29. The immunoglobulin according to any one of claims 24 to 26 which comprises an IgG, IgA, IgD or IgM sequence.
30. The immunoglobulin according to claim 20 which comprises an IgE complementarity determining region.
31. The immunoglobulin according to claim 30 which is an IgE analogue having a variant amino acid sequence within about residues 373 to 390, inclusive or residues 446 to 453, inclusive.

EP 1 260 521 A2

32. The immunoglobulin according to claims 30 which is an IgE analogue having a variant amino acid sequence within about residues 382 to 390, inclusive or residues 446 to 453, inclusive.

33. The immunoglobulin according to claim 32, further comprising a Fc ϵ RI-binding loop EF and beta strand D domain.

34. The immunoglobulin according to claim 32, further comprising an immunoglobulin variable domain capable of binding a predetermined antigen, an enzyme or diagnostic label.

35. The immunoglobulin according to claim 34, wherein the antigen is CD8 or CD3.

36. The immunoglobulin according to claim 34, wherein the antigen is a lymphoid cell surface antigen.

37. The immunoglobulin according to any one of claims 30 to 36 which comprises an IgG, IgA, IgD or IgM sequence.

38. The immunoglobulin according to any one of claims 30 to 37 which binds Fc ϵ RII with at least about 75% of the affinity of native IgE, and binds Fc ϵ RI with no greater than about 10% of the efficiency of native IgE.

39. An antibody according to any one of the preceding claims for use in a method of medical treatment.

40. Use of an antibody according to any of claims 1 to 38 for the preparation of a medicament for the treatment of an allergic disorder.

25 11 FIG. 1 5 12

β-strand F loop FG β-strand G

450 E G E T Y Q C R V T H P H L P R A L X M R S T K T S G P

MaE11 Light Chain

DIVLTQSPASLAVSLGQRATISCKASQSVDDYDGSYMNWYQQKPGQPPILLIYAASYLG
SEIPARFSGSGSGTDFTLNHPVEEEDAATFYCQQSHEDPYTFGAGTKLEIK

MaE11 Heavy Chain

DVQLQESGPGLVKPSQSLSLACSVTGYSITSGYSWNWIRQFPGNKLEWMGSITYDGSS
NYNPSLKNRISVTRDTSQNQFFLKLNSATAEDTATYYCARGSHYFGHWHFAVWGAGTTVT
VSSAKTTPPSVYPLAR

Mae13 Light Chain

DIVMTQSQKFMSTSVGDRVSVTCKASQNVSSNVAWYQQKPGQSPKALIYSASYRYSV
PDRFTGSGSGTDFTLTISNVQSEDLAEYFCQQYYTYPLYTFGGGTKLEIKRADAAPTVSI
FPPSTR

Mae13 Heavy Chain

DVQLQESGPGLVKPSQSLSLTCTVTGYTITSDNAWNWIRQFPGNKLEWMGYINHSGTT
SYNPSLKSRIISITRDTSKNQFFLQLNSVTTEDTATYYCAWVVAAYAMDYWGQGTSTVTVSSA
KTTTPPSVYPLAR

Mae15 Light Chain

DIQLTQSPASLAVSLGQRATISCKASQSVDDYDGSYMNWYQQKPGQPPKLLIYAASNLES
GIPARFSGSGSGTDFTLNHPVEEEDAATYYCQQSNBDPFTFGAGT

Mae15 Heavy Chain

DVQHQSEPDVLKPSQSLSLTCTVTGYTITSGYNRHWIRQFPGNKLEWMGYIHYSGST
NYNPSLKRRIISITRDTSKNQFFLQLNSVTTEDTATYYCARGSIYYYGSRYRYFDVWGAGT
TVTVSSAKRHPHLSIHWPG

FIG.2

Humanized MaE11 Version 1 (intact IgG)

Heavy Chain

EVQLVESGGGLVQPGGSLRLSCAIVSGYSITSGYSWNWIRQAPGKGLEWVASITYDGSTNY
ADSVKGRFTISRDDSKNTFYLMNSLRAEDTAVYYCARGSHYFGHWHFAVWGQGTLLTVS
SASTKGKGPSVFLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL
QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEL
LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE
QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS
REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFPLYSKLTVDK
SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Light Chain

DIQLTQSPSSLSASVGDRVTITCRASQSVDDYDGDSYMNWYQQKPGKAPKLLIYAASYLES
GVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQSHEDPYTFGQGTKVEIKRTVAAPSVF
IFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLS
STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

FIG.3




FIG.4a




FIG.4b




FIG.5a




FIG.5b





FIG.5c



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☒ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.